

jc598 U.S. PTO  
05/13/99

A Bof  
Seq.

**"EXPRESS MAIL CERTIFICATE"**

**"EXPRESS MAIL" MAILING LABEL NUMBER EL213569493US**

**DATE OF DEPOSIT MAY 13, 1999.**

**I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING DEPOSITED  
WITH THE UNITED STATES POSTAL SERVICE "EXPRESS MAIL POST  
OFFICE TO ADDRESSEE" SERVICE UNDER 37 CFR 1.10 ON THE DATE  
INDICATED ABOVE AND IS ADDRESSED TO ASSISTANT  
COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231**

**NAME OF PERSON MAILING PAPER OR FEE  
(TYPE OR PRINT) NANCY E. ROUTSON**

**SIGNATURE Nancy E. Routson**

jc542 U.S. PTO  
09/31/89  
05/13/99

Attorney Docket No.: 0571R2

Box PATENT APPLICATION  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Transmitted herewith for filing is the patent application of:

Inventor(s): Aragula Gururaj Rao and Keith R. Roesler

Title: PROTEINS WITH INCREASED LEVELS OF ESSENTIAL  
AMINO ACIDS

Dear Sir:

Enclosed are the following papers:

- ☒ Specification
- ☐ Formal drawings (\_\_\_\_\_ sheets).
- ☒ Informal drawings (2 sheets and 2 copies thereof = 4).
- ☐ Form PTO 1449 and copies of art cited.
- ☐ An assignment of the invention to Pioneer Hi-Bred International, Inc.  
Please record the assignment and return it to the undersigned attorney/agent.
- ☐ A certified copy of each of the following applications of which priority under 35  
USC 119 is claimed in the enclosed declaration.

09341689 "051399"

Attorney Docket No.: 0571R2

- ☒ Submission of sequence listing: computer readable diskette and statement to support sequence listing.
- ☒ The total fee is calculated as shown below:
- |                                                                    |                          |
|--------------------------------------------------------------------|--------------------------|
| Basic Filing fee                                                   | \$750.00                 |
| Total Claims 51-20 = 31 x \$22.00                                  | 682.00                   |
| Independent Claims 18 - 3 = 15 x \$78.00                           | 1,170.00                 |
| <input type="checkbox"/> Multiple Dependent Claim present \$250.00 |                          |
| <b>TOTAL FILING FEE</b>                                            | <b><u>\$2,602.00</u></b> |
- ☒ Charge **\$2,602.00** to Deposit Account No. 16-1852.  
Two duplicate copies of this form are enclosed.
- ☒ The Commissioner is hereby authorized to charge any additional fees under 37 CFR 1.16 or 1.17 which may be required by this paper, or credit any overpayment to our Deposit Account No. 16-1852.

Respectfully submitted,

*Catherine D Brooke*

Catherine D. Brooke  
Agent for Applicants  
Registration No. 44,041

PIONEER HI-BRED INTERNATIONAL, INC.  
Corporate Intellectual Property  
7100 N.W. 62<sup>nd</sup> Avenue  
P.O. Box 1000  
Johnston, Iowa 50131-1000  
Phone: (515) 248-4819  
Facsimile: (515) 334-6883

666750-6891660

## **PROTEINS WITH INCREASED LEVELS OF ESSENTIAL AMINO ACIDS**

**Aragula Gururaj Rao**

**Keith R. Roesler**

### **5 CROSS-REFERENCE TO RELATED APPLICATIONS:**

This application claims priority from and is a continuation-in-part of pending U.S. Patent Application No. 08/740,682 filed Nov. 1, 1996. This application also claims priority from and is a continuation-in-part of U.S. Patent Application No. 09/297,418 filed April 30, 1999, which claims priority from  
10 PCT/US97/20441, filed October 31, 1997.

U.S. Patent Application No. 08/740,682, U.S. Patent Application No. 09/297,418 and PCT/US97/20441 are incorporated by reference herewithin.

### **Field of the Invention**

15 The present invention relates to the field of protein engineering wherein changing amino acid compositions effects improvements in the nutrition content of feed and food. Specifically, the present invention relates to methods of enhancing the nutritional content of animal feed by expressing derivatives of a  
20 protease inhibitor to provide higher percentages of essential amino acids in plants.

### **Background of the Invention**

25 Feed formulations are required to provide animals essential nutrients critical to growth. However, crop plants are generally rendered food sources of poor nutritional quality because they contain low proportions of several amino acids which are essential for, but cannot be synthesized by, monogastric animals.

30 For many years researchers have attempted to improve the balance of essential amino acids in the seed proteins of important crops through breeding programs. As more becomes known about seed storage proteins and the expression of the genes which encode these proteins, and as transformation systems are developed for a greater variety of plants, molecular approaches for

improving the nutritional quality of seed proteins can provide alternatives to the more conventional approaches. Thus, specific amino acid levels can be enhanced in a given crop via biotechnology.

One alternative method is to express a heterologous protein of favorable amino acid composition at levels sufficient to obviate feed supplementation. For example, a number of seed proteins rich in sulfur amino acids have been identified. A key to good expression of such proteins involves efficient expression cassettes with tissue-preferred promoters. Not only must the gene-controlling regions direct the synthesis of high levels of mRNA, the mRNA must be translated into a stable protein and over-expression of this protein must not be detrimental to plant or animal health.

Among the essential amino acids needed for animal nutrition, often limiting in crop plants, are methionine, threonine, lysine, isoleucine, leucine, valine, tryptophan, phenylalanine, and histidine. Attempts to increase the levels of these free amino acids by breeding, mutant selection and/or changing the composition of the storage proteins accumulated in crop plants has met with limited success.

A transgenic example is the phaseolin-promoted Brazil nut 2S expression cassette. However, even though Brazil nut protein increases the amount of total methionine and bound methionine, thereby improving nutritional value, there appeared to be a threshold limitation as to the total amount of methionine that is accumulated in the seeds. The seeds remain insufficient as sources of methionine and methionine supplementation is required in diets utilizing soybeans.

An alternative to the enhancement of specific amino acid levels by altering the levels of proteins containing the desired amino acid is modification of amino acid biosynthesis. Recombinant DNA and gene transfer technologies have been applied to alter enzyme activity catalyzing key steps in the amino acid biosynthetic pathway. See Glassman, U.S. Patent No. 5,258,300; Galili, et al., European Patent Application No. 485970; (1992); incorporated herein in its entirety. However, modification of the amino acid levels in seeds is not always correlated with changes in the level of proteins that incorporate those amino acids. See Burrow, et al., Mol. Gen. Genet.; Vol. 241; pp. 431-439; (1993);

incorporated herein in its entirety by reference. Increases in free lysine levels in leaves and seeds have been obtained by selection for DHDPS mutants or by expressing the E. coli DHDPS in plants. However, since the level of free amino acids in seeds, in general, is only a minor fraction of the total amino acid content, these increases have been insufficient to significantly increase the total amino acid content of seed.

The lysC gene is a mutant bacterial aspartate kinase which is desensitized to feedback inhibition by lysine and threonine. Expression of this gene results in an increase in the level of lysine and threonine biosynthesis. However, expression of this gene with seed-specific expression cassettes has resulted in only a 6-7% increase in the level of total threonine or lysine in the seed. See Karchi, et al., The Plant J.; Vol. 3; pp. 721-7; (1993); incorporated herein in its entirety by reference. Thus, there is minimal impact on the nutritional value of seeds, and supplementation with essential amino acids is still required.

In another study (Falco et al., Biotechnology 13:577-582, 1995), manipulation of bacterial DHDPs and aspartate kinase did result in useful increases in free lysine and total seed lysine. However, abnormal accumulation of lysine catabolites was also observed suggesting that the free lysine pool was subject to catabolism.

Based on the foregoing, there exists a need for methods of increasing the levels of essential amino acids in seeds of plants. Previous approaches have led to insufficient increases in the levels of both free and bound amino acids and insignificant enhancement of the nutritional content of the feed.

### **Summary of the Invention**

It is the object of the present invention to provide nucleic acids and polypeptides relating to the enhancement of essential amino acids in plants.

It is another object of the present invention to provide antigenic fragments of the polypeptides of the present invention.

It is another object of the present invention to provide transgenic plants comprising the nucleic acids of the present invention.



It is an object of this invention to provide a CI-2-like polypeptide with increased stability.

It is an object of the present invention to provide methods for increasing the essential amino acid content of plants.

It is an object of the present invention to provide methods for increasing the nutritional value of a protein by altering a CI-2-like polypeptide to enhance its nutritional value by substituting essential amino acids at positions corresponding to 1, 8, 11, 17, 18, 19, 20, 22, 23, 31, 34, 38, 40, 41, 47, 49, 56, 58, 59, 60, 61, 62, 63, 65, 67, 69, 73, 75, 76, 78, 79, 81, 82, or combinations thereof.

### **Detailed Description of the Invention**

Choices of substitutions described herewithin are optionally grouped within parentheses and are separated by a semicolon. The native amino acid preceeds the position number using SEQ ID NO. 2 as a reference. The possible substitutions follow the residue number.

### **Figure listing**

Figure 1 - Comparison of modified BHL sequences

Figure 2 - CI-2-like sequences

1. Hordeum vulgare (A01293)
2. Hordeum vulgare (Y08625)
3. Zea mays (S37493)
4. Vicia faba (A21463)
5. Cucurbita maxima (S55591, S12897)
6. Canavalia lineata (JC2380)
7. Vigna angularis (JX0089)
8. Nicotiana tabacum (S33547)
9. Nicotiana sylvestris (A56555)
10. Sambucus nigra (Z46949)
11. Momordica charantia (JC2508)
12. Cucurbita maxima (S12897)
13. Solanum tuberosum (A01291, U30861)
14. Solanum tuberosum (U30861)
15. Lycopersicon peruvianum (A39547)
16. Lycopersicon esculentum (A32067, A24048)
17. Lycopersicon esculentum (A24048)
18. Amaranthus caudatus (S40496)
19. Arabidopsis thaliana (AC005770)

### **Sequence identification**

Full length wild-type chymotrypsin inhibitor (WT CI-2) is coded for by the polypeptides of SEQ ID No. 2 which is encoded for by the nucleic acid of SEQ ID No. 1.

Truncated wild-type chymotrypsin inhibitor (WT CI-2) is coded for by the polypeptides of SEQ ID No. 4 which is encoded for by the nucleic acid of SEQ ID No. 3.

Barley High Lysine 1(BHL1) is coded for by the polypeptides of SEQ ID No. 6 which is encoded for by the nucleic acid of SEQ ID No. 5.

Barley High Lysine 2 (BHL2) is coded for by the polypeptides of SEQ ID No. 8 which is encoded for by the nucleic acid of SEQ ID No. 7.

Barley High Lysine 3 (BHL3) is coded for by the polypeptides of SEQ ID No. 10 which is encoded for by the nucleic acid of SEQ ID No. 9.

Barley High Lysine 3N (BHL3N) is coded for by the polypeptides of SEQ ID No. 12 which is encoded for by the nucleic acid of SEQ ID No. 11.

Barley High Lysine 4 (BHL4) is coded for by the polypeptides of SEQ ID No. 14 which is encoded for by the nucleic acid of SEQ ID No. 13.

Barley High Lysine 5 (BHL5) is coded for by the polypeptides of SEQ ID No. 16 which is encoded for by the nucleic acid of SEQ ID No. 15.

Barley High Lysine 6 (BHL6) is coded for by the polypeptides of SEQ ID No. 8 which is encoded for by the nucleic acid of SEQ ID No. 17.

Barley High Lysine 8 (BHL8) is coded for by the polypeptides of SEQ ID No. 20 which is encoded for by the nucleic acid of SEQ ID No. 19.

The 5' and 3' PCR primer pairs A & B, are identified as SEQ ID Nos. 21 and 22, respectively.

Maize EST PI-1 is coded for by the polypeptides of SEQ ID No.24 which is encoded for by the nucleic acid of SEQ ID No. 23.

Maize EST PI-2 is coded for by the polypeptides of SEQ ID No.26 which is encoded for by the nucleic acid of SEQ ID No. 25.

Maize EST PI-3 is coded for by the polypeptides of SEQ ID No.28 which is encoded for by the nucleic acid of SEQ ID No. 27.



Maize EST PI-4 is coded for by the polypeptides of SEQ ID No. 30 which is encoded for by the nucleic acid of SEQ ID No. 29.

Maize EST PI-5 is coded for by the polypeptides of SEQ ID No. 32 which is encoded for by the nucleic acid of SEQ ID No. 31.

5

It has been unexpectedly discovered that one class of compounds, derivatives of chymotrypsin inhibitor-2 ("CI-2"), can be modified to enhance its essential amino acid content. In a preferred embodiment of the present invention, the CI-2 derivatives simultaneously exhibit both enhanced essential amino acids. The present compounds are thus excellent candidates for feed grain and food transformation to enhance nutrition.

10

## Definitions

Units, prefixes, and symbols may be denoted in their SI accepted form.

- 5 Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. Numeric ranges are inclusive of the numbers defining the range. Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes. The terms defined below are more fully defined by reference to the specification as a whole.

- 15 A "CI-2 derived" polypeptide refers to a chymotrypsin inhibitor polypeptide that may be truncated or modified, substituted or have an amino terminal extension or an insert.

- A "CI-2 like" polypeptide refers to a polypeptide of at least 23 consecutive amino acids of Seq. ID No. 2 or 4; or a polypeptide of at least 30% amino acid sequence identity with corresponding region of Seq. ID Nos. 2 or 4 or 20; or a CI-2-like polypeptide with modifications identified in CI-2; or a protease inhibitor with an active site loop typically between 53 and 70; or a CI-2 homologue modified to enhance its nutritional value by altering the amino acid residues at positions corresponding to those defined herein. The following organisms (Genbank Accession Numbers) may be modified according to the methods and figures in the specification Hordeum vulgare (A01293), Hordeum vulgare (Y08625), Zea mays (S37493), Vicia faba (A21463), Cucurbita maxima (S55591, S12897), Canavalia lineata (JC2380), Vigna angularis (JX0089), Nicotiana tabacum (S33547), Nicotiana glauca (A56555), Sambucus nigra (Z46949), Momordica charantia (JC2508), Cucurbita maxima (S12897), Solanum tuberosum (A01291, U30861), Solanum tuberosum (U30861), Lycopersicon peruvianum (A39547), Lycopersicon esculentum (A32067, A24048), Lycopersicon esculentum (A24048), Amaranthus caudatus (S40496), Arabidopsis thaliana (AC005770).
- 20
- 25
- 30

“Nutritionally-enhancing” refers to adding nutritional components that could include essential amino acids, fat, oil, and or vitamins and other compositions imparting characteristics desired in feed.

“%” refers to molar % unless otherwise specified or implied.

5 “Essential amino acids” are amino acids that must be obtained from an external source because they are not synthesized by the individual. They are: methionine, threonine, lysine, isoleucine, leucine, valine, tryptophan, phenylalanine, and histidine.

10 The term "antibody" includes reference to antigen binding forms of antibodies (e.g., Fab, F(ab)<sub>2</sub>). The term "antibody" refers to a polypeptide substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof which specifically bind and recognize an analyte (antigen). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be  
15 synthesized *de novo* either chemically or by utilizing recombinant DNA methodology.

The term "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode  
20 identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position  
25 where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations" and represent one species of conservatively modified variation. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the  
30 nucleic acid. One of ordinary skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally

identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide of the present invention is implicit in each described polypeptide sequence and incorporated herein by reference.

As to amino acid sequences, one of skill will recognize that individual  
5 substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Thus, any number of amino acid residues selected  
10 from the group of integers consisting of from 1 to 15 can be so altered. Thus, for example, 1, 2, 3, 4, 5, 7, or 10 alterations can be made. Conservatively modified variants typically provide similar biological activity as the unmodified polypeptide sequence from which they are derived. For example, substrate specificity, enzyme activity, or ligand/receptor binding is generally at least 30%, 40%, 50%,  
15 60%, 70%, 80%, or 90% of the native protein for its native substrate. Conservative substitution tables providing functionally similar amino acids are well known in the art.

The following six groups each contain amino acids that are conservative substitutions for one another:

- 20 1) Alanine (A), Serine (S), Threonine (T), Cysteine (C);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 25 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

See also, Creighton (1984) Proteins W.H. Freeman and Company.

The following groups each contain amino acids that are conservative and essential amino acid substitutions for one another:

- 1) Threonine (T), and Lysine (K)
- 30 2) Isoleucine (I), Leucine (L), Methionine (M), and Valine (V).

The term "isolated" refers to material, such as a nucleic acid or a protein, which is: (1) substantially or essentially free from components which normally accompany or interact with the material as found in its naturally occurring environment or (2) if the material is in its natural environment, the material has  
 5 been altered by deliberate human intervention to a composition and/or placed at a locus in the cell other than the locus native to the material.

As used herein, "polypeptide" means proteins, protein fragments, modified proteins, amino acid sequences and synthetic amino acid sequences. The polypeptide can be glycosylated or not.

10 As used herein, "plant" includes but is not limited to plant cells, plant tissue and plant seeds.

As used herein, "promoter" includes reference to a region of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription.

15 By "fragment" is intended a portion of the nucleotide sequence or a portion of the amino acid sequence and hence protein encoded thereby. Preferably fragments of a nucleotide sequence may encode protein fragments that retain the biological activity of the native nucleic acid. However, fragments of a nucleotide sequence which are useful as hybridization probes generally do not encode  
 20 fragment proteins retaining biological activity. Fragments of a nucleotide sequence are generally greater than 10 nucleotides, preferably at least 20 nucleotides and up to the entire nucleotide sequence encoding the proteins of the invention. Generally probes are less than 1000 nucleotides and preferably less than 500 nucleotides. Fragments of the invention include antisense sequences  
 25 used to decrease expression of the inventive nucleic acids. Such antisense fragments may vary in length ranging from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, up to and including the entire coding sequence.

30 By "variants" is intended substantially similar sequences. Generally, nucleic acid sequence variants of the invention will have at least 40%, 50%, 55%, 60%, 70%, or preferably 80%, more preferably at least 90% and most preferably at least 95% sequence identity to the native nucleotide sequence.

Generally, polypeptide sequence variants of the invention will have at least about 55%, 60%, 70%, 80%, or preferably at least about 90% and more preferably at least about 95% sequence identity to the modified protein.

As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences includes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. A polypeptide is substantially identical to a second polypeptide, for example, where the two polypeptides differ only by conservative substitution.

Methods of alignment of sequences for comparison are well-known in the art. For purposes of defining the present invention, the BLAST 2.0 suite of programs using default parameters is used. Altschul *et al.*, *Nucleic Acids Res.* 25:3389-3402 (1997). Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>).

By "functionally equivalent" is intended that the sequence of the variant defines a chain that produces a protein having substantially the same biological effect as the native protein of interest.

When the nucleic acid is prepared or altered synthetically, advantage can be taken of known codon preferences of the intended host where the nucleic acid is to be expressed. For example, although nucleic acid sequences of the present invention may be expressed in both monocotyledonous and dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* *Nucl. Acids Res.* 17: 477-498 (1989)). Thus, the maize preferred codon for a particular amino acid may be derived from known gene sequences from maize. Maize codon usage for 28 genes from maize plants are listed in Table 4 of Murray *et al.*, *supra*.

By "immunologically reactive conditions" is meant conditions which allow an antibody, generated to a particular epitope, to bind to that epitope to a

detectably greater degree (e.g., at least 2-fold over background) than the antibody binds to substantially all other epitopes. Immunologically reactive conditions are dependent upon the format of the antibody binding reaction and typically are those utilized in immunoassay protocols. See Harlow and Lane, 5 *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York (1988), for a description of immunoassay formats and conditions.

The terms "isolated" or "biologically pure" refer to material which is: (1) substantially or essentially free from components which normally accompany or interact with it as found in its naturally occurring environment. The isolated 10 material optionally comprises material not found with the material in its natural environment. (2) If the material is in its natural environment, the material has been synthetically (non-naturally) altered to a composition and/or placed at a locus in the cell (e.g., genome) not native to a material found in that environment. The alteration to yield the synthetic material can be performed on the material 15 within or removed from its natural state. For example, a naturally occurring nucleic acid becomes an isolated nucleic acid if it is altered, or if it is transcribed from DNA which is altered, by non-natural, synthetic (i.e., "man-made") methods performed within the cell from which it originates. See, e.g., Compounds and Methods for Site Directed Mutagenesis in Eukaryotic Cells, Kmiec, U.S. Patent 20 No. 5,565,350; *In Vivo* Homologous Sequence Targeting in Eukaryotic Cells; Zarling *et al.*, PCT/US93/03868. Likewise, a naturally occurring nucleic acid (e.g., a promoter) become isolated if it is introduced by non-naturally occurring means to a locus of the genome not native to that nucleic acid.

As used herein "operably linked" includes reference to a functional linkage 25 between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

30 As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds and plant cells and progeny of same. Plant cell, as used herein includes, without limitation, seeds, suspension





demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. Such modifications are well known to those of skill and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as, for instance *Proteins - Structure and Molecular Properties*, 2nd ed., T. E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as, for example, those provided by Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pp. 1-12 in *Posttranslational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter *et al.*, Meth. Enzymol. 182: 626-646 (1990) and Rattan *et al.*, *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). It will be appreciated, as is well known and as noted above, that polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of posttranslation events, including natural processing event and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translation natural process and by entirely synthetic methods, as well. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention, as well. For instance, the amino terminal residue of polypeptides made in *E. coli* or other cells, prior to proteolytic processing, almost invariably will be N-formylmethionine. During post-translational modification of the peptide, a methionine residue at the NH<sub>2</sub>-terminus may be deleted.

Accordingly, this invention contemplates the use of both the methionine-containing and the methionineless amino terminal variants of the protein of the invention. In general, as used herein, the term polypeptide encompasses all such modifications, particularly those that are present in polypeptides synthesized by  
5 expressing a polynucleotide in a host cell.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using one of the alignment  
10 programs described using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means  
15 sequence identity of at least 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes.

## 20 **NUCLEIC ACIDS**

The isolated nucleic acids of the present invention can be made using (a) standard recombinant methods, (b) synthetic techniques, or combinations thereof. In some embodiments, the polynucleotides of the present invention will be cloned, amplified, or otherwise constructed from a monocot or dicot. In preferred  
25 embodiments the monocot is corn, sorghum, barley, wheat, millet, or rice. Preferred dicots include soybeans, sunflower, canola, alfalfa, cotton, potato, lupin or cassava.

Functional fragments included in the invention can be obtained using primers that selectively hybridize under stringent conditions. Primers are  
30 generally at least 12 bases in length and can be as high as 200 bases, but will generally be from 15 to 75, preferably from 15 to 50. Functional fragments can

be identified using a variety of techniques such as restriction analysis, Southern analysis, primer extension analysis, and DNA sequence analysis.

The present invention includes a plurality of polynucleotides that encode for the identical amino acid sequence. The degeneracy of the genetic code  
5 allows for such "silent variations" which can be used, for example, to selectively hybridize and detect allelic variants of polynucleotides of the present invention. Additionally, the present invention includes isolated nucleic acids comprising allelic variants. The term "allele" as used herein refers to a related nucleic acid of the same gene.

10 Variants of nucleic acids included in the invention can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like. See, for example, Ausubel, pages 8.0.3 - 8.5.9. Also, see generally, McPherson (ed.), *DIRECTED MUTAGENESIS: A Practical approach*, (IRL Press, 1991). Thus, the  
15 present invention also encompasses DNA molecules comprising nucleotide sequences that have substantial sequence similarity with the inventive sequences.

Variants included in the invention may contain individual substitutions, deletions or additions to the nucleic acid or polypeptide sequences. Such  
20 changes will alter, add or delete a single amino acid or a small percentage of amino acids in the encoded sequence. Variants are referred to as "conservatively modified variants" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. When the nucleic acid is prepared or altered synthetically, advantage can be taken of known codon  
25 preferences of the intended host.

The present invention also includes the use of 5' and/or 3' UTR regions for modulation of translation of heterologous coding sequences. Positive sequence motifs include translational initiation consensus sequences (Kozak, *Nucleic Acids Res.* 15:8125 (1987)) and the 7-methylguanosine cap structure (Drummond *et al.*,  
30 *Nucleic Acids Res.* 13:7375 (1985)). Negative elements include stable intramolecular 5' UTR stem-loop structures (Muesing *et al.*, *Cell* 48:691 (1987))

and AUG sequences or short open reading frames preceded by an appropriate AUG in the 5' UTR (Kozak, *supra*, Rao *et al.*, *Mol. and Cell. Biol.* 8:284 (1988)).

Further, the polypeptide-encoding segments of the polynucleotides of the present invention can be modified to alter codon usage. Altered codon usage  
5 can be employed to alter translational efficiency and/or to optimize the coding sequence for expression in a desired host or to optimize the codon usage in a heterologous sequence for expression in maize. Codon usage in the coding regions of the polynucleotides of the present invention can be analyzed statistically using commercially available software packages such as "Codon  
10 Preference" available from the University of Wisconsin Genetics Computer Group (see Devereaux *et al.*, *Nucleic Acids Res.* 12: 387-395 (1984)) or MacVector 4.1 (Eastman Kodak Co., New Haven, Conn.).

For example, the inventive nucleic acids can be optimized for enhanced expression in organisms of interest. See, for example, EPA0359472;  
15 WO91/16432; Perlak *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:3324-3328; and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498. In this manner, the genes can be synthesized utilizing species-preferred codons. See, for example, Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, the disclosure of which is incorporated herein by reference.

20 The present invention provides subsequences comprising isolated nucleic acids containing at least 16 contiguous bases of the inventive sequences. For example the isolated nucleic acid includes those comprising at least 20, 25, 30, 40, 50, 60, 75 or 100 contiguous nucleotides of the inventive sequences. Subsequences of the isolated nucleic acid can be used to modulate or detect  
25 gene expression by introducing into the subsequences compounds which bind, intercalate, cleave and/or crosslink to nucleic acids.

The nucleic acids of the invention may conveniently comprise a multi-cloning site comprising one or more endonuclease restriction sites inserted into the nucleic acid to aid in isolation of the polynucleotide. Also, translatable  
30 sequences may be inserted to aid in the isolation of the translated polynucleotide of the present invention. For example, a hexa-histidine marker sequence provides a convenient means to purify the proteins of the present invention.

A polynucleotide of the present invention can be attached to a vector, adapter, promoter, transit peptide or linker for cloning and/or expression of a polynucleotide of the present invention. Additional sequences may be added to such cloning and/or expression sequences to optimize their function in cloning and/or expression, to aid in isolation of the polynucleotide, or to improve the introduction of the polynucleotide into a cell. Use of cloning vectors, expression vectors, adapters, and linkers is well known and extensively described in the art. For a description of such nucleic acids see, for example, Stratagene Cloning Systems, Catalogs 1995, 1996, 1997 (La Jolla, CA); and, Amersham Life Sciences, Inc, Catalog '97 (Arlington Heights, IL).

The isolated nucleic acid compositions of this invention, such as RNA, cDNA, genomic DNA, or a hybrid thereof, can be obtained from plant biological sources using any number of cloning methodologies known to those of skill in the art. In some embodiments, oligonucleotide probes which selectively hybridize, under stringent conditions, to the polynucleotides of the present invention are used to identify the desired sequence in a cDNA or genomic DNA library.

Exemplary total RNA and mRNA isolation protocols are described in *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997); and, *Current Protocols in Molecular Biology*, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Total RNA and mRNA isolation kits are commercially available from vendors such as Stratagene (La Jolla, CA), Clontech (Palo Alto, CA), Pharmacia (Piscataway, NJ), and 5'-3' (Paoli, PA). See also, U.S. Patent Nos. 5,614,391; and, 5,459,253.

Typical cDNA synthesis protocols are well known to the skilled artisan and are described in such standard references as: *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997); and, *Current Protocols in Molecular Biology*, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). cDNA synthesis kits are available from a variety of commercial vendors such as Stratagene or Pharmacia.

Typically, stringent hybridization conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least

about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

Preferably the hybridization is conducted under low stringency conditions which include hybridization with a buffer solution of 30 % formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulfate) at 37°C for 24 hrs., and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50°C. More preferably the hybridization is conducted under moderate stringency conditions which include hybridization in 40 % formamide, 1 M NaCl, 1% SDS at 37°C for 24 hrs., and a wash in 0.5X to 1X SSC at 55°C. Most preferably the hybridization is conducted under high stringency conditions which include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C for 24 hrs., and a wash in 0.1X SSC at 60°C.

An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New York (1993); and *Current Protocols in Molecular Biology*, Chapter 2, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995). Often, cDNA libraries will be normalized to increase the representation of relatively rare cDNAs.

The nucleic acids of the invention can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of polynucleotides of the present invention and related genes directly from genomic DNA or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Examples of techniques useful for *in vitro* amplification methods are found in Berger, Sambrook, and Ausubel, as well as Mullis *et al.*, U.S. Patent No. 4,683,202 (1987); and, *PCR Protocols A Guide to Methods and Applications*, Innis *et al.*, Eds., Academic Press Inc., San Diego, CA (1990). Commercially

available kits for genomic PCR amplification are known in the art. See, e.g., Advantage-GC Genomic PCR Kit (Clontech). The T4 gene 32 protein (Boehringer Mannheim) can be used to improve yield of long PCR products.

PCR-based screening methods have also been described. Wilfinger *et al.* describe a PCR-based method in which the longest cDNA is identified in the first step so that incomplete clones can be eliminated from study. *BioTechniques*, 22(3): 481-486 (1997).

In one aspect of the invention, nucleic acids can be amplified from a *Zea mays* nucleic acid library. The nucleic acid library may be a cDNA library, a genomic library, or a library generally constructed from nuclear transcripts at any stage of intron processing.

Libraries can be made from a variety of maize tissues. Good results have been obtained using mitotically active tissues such as shoot meristems, shoot meristem cultures, embryos, callus and suspension cultures, immature ears and tassels, and young seedlings. The cDNA of the present invention was obtained from developing endosperm. Since cell cycle proteins are typically expressed at specific cell cycle stages it may be possible to enrich for such rare messages using exemplary cell cycle inhibitors such as aphidicolin, hydroxyurea, mimosine, and double-phosphate starvation methods to block cells at the G1/S boundary. Cells can also be blocked at this stage using the double phosphate starvation method. Hormone treatments that stimulate cell division, for example cytokinin, would also increase expression of the cell cycle RNA.

Alternatively, the sequences of the invention can be used to isolate corresponding sequences in other organisms, particularly other plants, more particularly, other monocots. In this manner, methods such as PCR, hybridization, and the like can be used to identify such sequences having substantial sequence similarity to the sequences of the invention. See, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). and Innis *et al.* (1990), *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York). Coding sequences isolated based on their sequence identity to the

entire inventive coding sequences set forth herein or to fragments thereof are encompassed by the present invention.

The isolated nucleic acids of the present invention can also be prepared by direct chemical synthesis by methods such as the phosphotriester method of Narang *et al.*, *Meth. Enzymol.* 68: 90-99 (1979); the phosphodiester method of Brown *et al.*, *Meth. Enzymol.* 68: 109-151 (1979); the diethylphosphoramidite method of Beaucage *et al.*, *Tetra. Lett.* 22: 1859-1862 (1981); the solid phase phosphoramidite triester method described by Beaucage and Caruthers, *Tetra. Letts.* 22(20): 1859-1862 (1981), *e.g.*, using an automated synthesizer, *e.g.*, as described in Needham-VanDevanter *et al.*, *Nucleic Acids Res.*, 12: 6159-6168 (1984); and, the solid support method of U.S. Patent No. 4,458,066. Chemical synthesis generally produces a single stranded oligonucleotide. This may be converted into double stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill will recognize that while chemical synthesis of DNA is limited to sequences of about 100 bases, longer sequences may be obtained by the ligation of shorter sequences.

### **EXPRESSION CASSETTES**

In another embodiment expression cassettes comprising isolated nucleic acids of the present invention are provided. An expression cassette will typically comprise a polynucleotide of the present invention operably linked to transcriptional initiation regulatory sequences which will direct the transcription of the polynucleotide in the intended host cell, such as tissues of a transformed plant.

The construction of expression cassettes that can be employed in conjunction with the present invention is well known to those of skill in the art in light of the present disclosure. See, *e.g.*, Sambrook, *et al.*; Molecular Cloning: A Laboratory Manual; Cold Spring Harbor, New York; (1989); Gelvin, *et al.*; Plant Molecular Biology Manual; (1990); Plant Biotechnology: Commercial Prospects and Problems, eds. Prakash, *et al.*; Oxford & IBH Publishing Co.; New Delhi, India; (1993); and Heslot, *et al.*; Molecular Biology and Genetic Engineering of



Yeasts; CRC Press, Inc., USA; (1992); each incorporated herein in its entirety by reference.

For example, plant expression vectors may include (1) a cloned plant nucleic acid under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (e.g., one conferring inducible, constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific/selective expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

Constitutive, tissue-preferred or inducible promoters can be employed. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (U.S. Patent No. 5,683,439), the *Nos* promoter, the pEmu promoter, the rubisco promoter, the GRP1-8 promoter and other transcription initiation regions from various plant genes known to those of skill.

An efficient plant promoter that may be used is an overproducing plant promoter. Overproducing plant promoters that may be used in this invention include the promoter of the chlorophyll  $\alpha$ - $\beta$  binding protein, and the promoter of the small sub-unit (ss) of the ribulose-1,5-biphosphate carboxylase from soybean. See e.g. Berry-Lowe, et al., J. Molecular and App. Gen.; Vol. 1; pp. 483-498; (1982); incorporated herein in its entirety by reference. These two promoters are known to be light-induced, in eukaryotic plant cells. See e.g., An Agricultural Perspective, A. Cashmore, Pelham, New York, 1983, pp. 29-38, G. Coruzzi, et al., J. Biol. Chem., Vol. 258; p. 1399 (1983), and P. Dunsmuir, et al., J. Molecular and App. Gen., Vol. 2; p. 285 (1983); all incorporated herein in their entirety by reference.

Examples of inducible promoters are the Adh1 promoter which is inducible by hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress,

and the PPKK promoter which is inducible by light. Also useful are promoters which are chemically inducible.

Examples of promoters under developmental control include promoters that initiate transcription preferentially in certain tissues, such as leaves, roots, fruit, seeds, or flowers. An exemplary promoter is the anther specific promoter 5126 (U.S. Patent Nos. 5,689,049 and 5,689,051). Examples of seed-preferred promoters include, but are not limited to, 27 kD gamma zein promoter and waxy promoter, Boronat,A., Martinez,M.C., Reina,M., Puigdomenech,P. and Palau,J.; Isolation and sequencing of a 28 kD glutelin-2 gene from maize: Common elements in the 5' flanking regions among zein and glutelin genes; Plant Sci. 47, 95-102 (1986) and Reina,M., Ponte,I., Guillen,P., Boronat,A. and Palau,J., Sequence analysis of a genomic clone encoding a Zc2 protein from Zea mays W64 A, Nucleic Acids Res. 18 (21), 6426 (1990). See the following site relating to the waxy promoter: Kloesgen,R.B., Gierl,A., Schwarz-Sommer,ZS. and Saedler,H., Molecular analysis of the waxy locus of Zea mays, Mol. Gen. Genet. 203, 237-244 (1986). Promoters that express in the embryo, pericarp, and endosperm are disclosed in US applications Ser. Nos. 60/097,233 filed August 20, 1998 and 60/098,230 filed August 28, 1998. The disclosures each of these are incorporated herein by reference in their entirety.

Either heterologous or non-heterologous (i.e., endogenous) promoters can be employed to direct expression of the nucleic acids of the present invention. These promoters can also be used, for example, in expression cassettes to drive expression of antisense nucleic acids to reduce, increase, or alter concentration and/or composition of the proteins of the present invention in a desired tissue.

If polypeptide expression is desired, it is generally desirable to include a polyadenylation region at the 3'-end of a polynucleotide coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA. The 3' end sequence to be added can be derived from, for example, the nopaline synthase or octopine synthase genes, or alternatively from another plant gene, or less preferably from any other eukaryotic gene.

An intron sequence can be added to the 5' untranslated region or the coding sequence of the partial coding sequence to increase the amount of the mature message that accumulates. See for example Buchman and Berg, *Mol. Cell Biol.* 8: 4395-4405 (1988); Callis *et al.*, *Genes Dev.* 1: 1183-1200 (1987).

5 Use of maize introns Adh1-S intron 1, 2, and 6, the Bronze-1 intron are known in the art. See generally, *The Maize Handbook*, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994).

The vector comprising the sequences from a polynucleotide of the present invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. Usually, the selectable marker gene will encode antibiotic or herbicide resistance. Suitable genes include those coding for resistance to the antibiotic spectinomycin or streptomycin (e.g., the *aadA* gene), the streptomycin phosphotransferase (SPT) gene coding for streptomycin resistance, the neomycin phosphotransferase (NPTII) gene encoding kanamycin or geneticin resistance, the hygromycin phosphotransferase (HPT) gene coding for hygromycin resistance.

Suitable genes coding for resistance to herbicides include those which act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonylurea-type herbicides (e.g., the acetolactate synthase (ALS) gene containing mutations leading to such resistance in particular the S4 and/or Hra mutations), those which act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the *bar* gene), or other such genes known in the art. The *bar* gene encodes resistance to the herbicide basta and the ALS gene encodes resistance to the herbicide chlorsulfuron.

25 Typical vectors useful for expression of nucleic acids in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.*, *Meth. In Enzymol.*, 153:253-277 (1987). Exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl *et al.*, *Gene*, 61:1-11 (1987) and Berger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101.2 that is available from Clontech Laboratories, Inc. (Palo Alto, CA).

A variety of plant viruses that can be employed as vectors are known in the art and include cauliflower mosaic virus (CaMV), geminivirus, brome mosaic virus, and tobacco mosaic virus.

## 5 **PROTEINS**

Proteins of the present invention include proteins derived from the native protein by deletion (so-called truncation), addition, or substitution of one or more amino acids at one or more sites in the native protein. Methods for such deletions, additions and substitutions are generally known in the art.

10 For example, amino acid sequence variants of the polypeptide can be prepared by mutations in the cloned DNA sequence encoding the native protein of interest. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York);  
15 Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods Enzymol.* 154:367-382; Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor, New York); U.S. Patent No. 4,873,192; and the references cited therein; herein incorporated by reference.

20 In constructing variants of the proteins of interest, modifications to the nucleotide sequences encoding the variants will be made such that variants continue to possess the desired activity. Obviously, any mutations made in the DNA encoding the variant protein must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See EP Patent Application Publication No. 75,444.

25 The isolated proteins of the present invention include a polypeptide comprising at least 23 contiguous amino acids encoded by any one of the nucleic acids of the present invention, or polypeptides which are conservatively modified variants thereof. The proteins of the present invention or variants thereof can comprise any number of contiguous amino acid residues from a polypeptide of  
30 the present invention, wherein that number is selected from the group of integers consisting of from 23 to the number of residues in a full-length polypeptide of the present invention. Optionally, this subsequence of contiguous amino acids is at

least 25, 30, 35, or 40 amino acids in length, often at least 50, 60, 70, 80, or 90 amino acids in length.

The present invention includes modifications that can be made to an inventive protein to increase nutritional enhancement activity. Some  
5 modifications may be made to facilitate the cloning, expression, or incorporation of the targeting molecule into a fusion protein. Such modifications are well known to those of skill in the art and include, for example, a methionine added at the amino terminus to provide an initiation site, or additional amino acids (e.g., poly His) placed on either terminus to create conveniently located restriction sites or  
10 termination codons or purification sequences.

A protein of the present invention can be expressed in a recombinantly engineered cell such as bacteria, yeast, insect, mammalian, or preferably plant cells. The cells produce the protein in a non-natural condition (e.g., in quantity, composition, location, and/or time), because they have been genetically altered  
15 through human intervention to do so.

Typically, an intermediate host cell will be used in the practice of this invention to increase the copy number of the cloning vector. With an increased copy number, the vector containing the nucleic acid of interest can be isolated in significant quantities for introduction into the desired plant cells.

20 Host cells that can be used in the practice of this invention include prokaryotes, including bacterial hosts such as *Eschericia coli*, *Salmonella typhimurium*, and *Serratia marcescens*. Eukaryotic hosts such as yeast or filamentous fungi may also be used in this invention. It is preferred to use plant promoters that do not cause expression of the polypeptide in bacteria.

25 Commonly used prokaryotic control sequences include promoters such as the beta lactamase (penicillinase) and lactose (lac) promoter systems (Chang *et al.*, Nature 198:1056 (1977)), the tryptophan (trp) promoter system (Goeddel *et al.*, Nucleic Acids Res. 8:4057 (1980)) and the lambda derived P L promoter and N-gene ribosome binding site (Shimatake *et al.*, Nature 292:128 (1981)). The  
30 inclusion of selection markers in DNA vectors transfected in *E. coli* is also useful. Examples of such markers include genes specifying resistance to ampicillin, tetracycline, or chloramphenicol.

The vector is selected to allow introduction into the appropriate host cell. Bacterial vectors are typically of plasmid or phage origin. Expression systems for expressing a protein of the present invention are available using *Bacillus sp.* and *Salmonella* (Palva, *et al.*, *Gene* 22: 229-235 (1983); Mosbach, *et al.*, *Nature* 302: 543-545 (1983)).

Synthesis of heterologous proteins in yeast is well known. See Sherman, F., *et al.*, *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory (1982). Two widely utilized yeast for production of eukaryotic proteins are *Saccharomyces cerevisiae* and *Pichia pastoris*. Vectors, strains, and protocols for expression in *Saccharomyces* and *Pichia* are known in the art and available from commercial suppliers (e.g., Invitrogen). Suitable vectors usually have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or alcohol oxidase, and an origin of replication, termination sequences and the like as desired.

A protein of the present invention, once expressed, can be isolated from yeast by lysing the cells and applying standard protein isolation techniques to the lysates. The monitoring of the purification process can be accomplished by using Western blot techniques or radioimmunoassay of other standard immunoassay techniques.

The proteins of the present invention can also be constructed using non-cellular synthetic methods. Solid phase synthesis of proteins of less than about 50 amino acids in length may be accomplished by attaching the C-terminal amino acid of the sequence to an insoluble support followed by sequential addition of the remaining amino acids in the sequence. Techniques for solid phase synthesis are described by Barany and Merrifield, *Solid-Phase Peptide Synthesis*, pp. 3-284 in *The Peptides: Analysis, Synthesis, Biology*. Vol. 2: *Special Methods in Peptide Synthesis, Part A.*; Merrifield, *et al.*, *J. Am. Chem. Soc.* 85: 2149-2156 (1963), and Stewart *et al.*, *Solid Phase Peptide Synthesis*, 2nd ed., Pierce Chem. Co., Rockford, Ill. (1984). Proteins of greater length may be synthesized by condensation of the amino and carboxy termini of shorter fragments. Methods of forming peptide bonds by activation of a carboxy terminal

end (e.g., by the use of the coupling reagent N,N'-dicyclohexylcarbodiimide) is known to those of skill.

The proteins of this invention may be purified to substantial purity by standard techniques well known in the art, including detergent solubilization, selective precipitation with such substances as ammonium sulfate, column chromatography, immunopurification methods, and others. See, for instance, R. Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag: New York (1982); Deutscher, *Guide to Protein Purification*, Academic Press (1990). For example, antibodies may be raised to the proteins as described herein. Purification from *E. coli* can be achieved following procedures described in U.S. Patent No. 4,511,503. Detection of the expressed protein is achieved by methods known in the art and include, for example, radioimmunoassays, Western blotting techniques or immunoprecipitation.

The present invention further provides a method for modulating (i.e., increasing or decreasing) the concentration or composition of the polypeptides of the present invention in a plant or part thereof. Modulation of the polypeptides can be effected by increasing or decreasing the concentration and/or the composition of the polypeptides in a plant. The method comprises transforming a plant cell with an expression cassette comprising a polynucleotide of the present invention to obtain a transformed plant cell, growing the transformed plant cell under plant forming conditions, and inducing expression of the polynucleotide in the plant for a time sufficient to modulate concentration and/or composition of the polypeptides in the plant or plant part.

In some embodiments, the content and/or composition of polypeptides of the present invention in a plant may be modulated by altering, *in vivo* or *in vitro*, the promoter of a non-isolated gene of the present invention to up- or down-regulate gene expression. In some embodiments, the coding regions of native genes of the present invention can be altered via substitution, addition, insertion, or deletion. See, e.g., Kmiec, U.S. Patent 5,565,350; Zarling *et al.*, PCT/US93/03868.

In some embodiments, an isolated nucleic acid (e.g., a vector) comprising a promoter sequence is transfected into a plant cell. Subsequently, a plant cell

comprising the isolated nucleic acid is selected for by means known to those of skill in the art such as, but not limited to, Southern blot, DNA sequencing, or PCR analysis using primers specific to the promoter and to the nucleic acid and detecting amplicons produced therefrom. A plant or plant part altered or modified by the foregoing embodiments is grown under plant forming conditions for a time sufficient to modulate the concentration and/or composition of polypeptides of the present invention in the plant. Plant forming conditions are well known in the art.

In general, concentration of the polypeptides is increased or decreased by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% relative to a native control plant, plant part, or cell lacking the aforementioned expression cassette. Modulation in the present invention may occur during and/or subsequent to growth of the plant to the desired stage of development.

Modulating nucleic acid expression temporally and/or in particular tissues can be controlled by employing the appropriate promoter operably linked to a polynucleotide of the present invention in, for example, sense or antisense orientation as discussed in greater detail above. Induction of expression of a polynucleotide of the present invention can also be controlled by exogenous administration of an effective amount of inducing compound. Inducible promoters and inducing compounds that activate expression from these promoters are well known in the art.

In preferred embodiments, the polypeptides of the present invention are modulated in monocots or dicots, preferably corn, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, millet, and lupin.

Means of detecting the proteins of the present invention are not critical aspects of the present invention. In a preferred embodiment, the proteins are detected and/or quantified using any of a number of well recognized immunological binding assays (see, e.g., U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, see also *Methods in Cell Biology*, Vol. 37: *Antibodies in Cell Biology*, Asai, Ed., Academic Press, Inc. New York (1993); *Basic and Clinical Immunology* 7th Edition, Stites & Terr, Eds. (1991). Moreover, the immunoassays of the present invention can be performed in any of several configurations, e.g., those reviewed in *Enzyme*



*Immunoassay*, Maggio, Ed., CRC Press, Boca Raton, Florida (1980); Tijan, Practice and Theory of Enzyme Immunoassays, *Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers B.V., Amsterdam (1985); Harlow and Lane, *supra*; *Immunoassay: A Practical Guide*, Chan, Ed., Academic Press, Orlando, FL (1987); *Principles and Practice of Immunoassays*, Price and Newman Eds., Stockton Press, NY (1991); and *Non-isotopic Immunoassays*, Ngo, Ed., Plenum Press, NY (1988).

Typical methods for detecting proteins include Western blot (immunoblot) analysis, analytic biochemical methods such as electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, and the like, and various immunological methods such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, and the like.

For a review of various labeling or signal producing systems which may be used, see, U.S. Patent No. 4,391,904, which is incorporated herein by reference.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies.

The proteins of the present invention can be used for identifying compounds that bind to (e.g., substrates), and/or increase or decrease (i.e., modulate) the activity of, catalytically active polypeptides of the present invention. The method comprises contacting a polypeptide of the present invention with a compound whose ability to bind to or modulate activity is to be determined.

Methods of measuring enzyme kinetics are well known in the art. See, e.g., Segel, *Biochemical Calculations*, 2<sup>nd</sup> ed., John Wiley and Sons, New York (1976).

Antibodies can be raised to a protein of the present invention, including individual, allelic, strain, or species variants, and fragments thereof, both in their naturally occurring (full-length) forms and in recombinant forms. Additionally, antibodies are raised to these proteins in either their native configurations or in non-native configurations. Anti-idiotypic antibodies can also be generated. Many methods of making antibodies are known to persons of skill. Description of

techniques for preparing such monoclonal antibodies are found in, e.g., *Basic and Clinical Immunology*, 4th ed., Stites *et al.*, Eds., Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane, *Supra*; Goding, *Monoclonal Antibodies: Principles and Practice*, 2nd ed., Academic Press, New York, NY (1986); and Kohler and Milstein, *Nature* 256: 495-497 (1975).

Other suitable techniques involve selection of libraries of recombinant antibodies in phage or similar vectors (see, e.g., Huse *et al.*, *Science* 246: 1275-1281 (1989); and Ward, *et al.*, *Nature* 341: 544-546 (1989); and Vaughan *et al.*, *Nature Biotechnology*, 14: 309-314 (1996)). Alternatively, high avidity human monoclonal antibodies can be obtained from transgenic mice comprising fragments of the unrearranged human heavy and light chain Ig loci (i.e., minilocus transgenic mice). Fishwild *et al.*, *Nature Biotech.*, 14: 845-851 (1996). Also, recombinant immunoglobulins may be produced. See, Cabilly, U.S. Patent No. 4,816,567; and Queen *et al.*, *Proc. Nat'l Acad. Sci.* 86: 10029-10033 (1989).

The antibodies of this invention can be used for affinity chromatography in isolating proteins of the present invention, for screening expression libraries for particular expression products such as normal or abnormal protein or for raising anti-idiotypic antibodies which are useful for detecting or diagnosing various pathological conditions related to the presence of the respective antigens.

Frequently, the proteins and antibodies of the present invention will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionucleotides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like.

### **Transfection/Transformation of Cells**

The method of transformation/transfection is not critical to the invention; various methods of transformation or transfection are currently available. As newer methods are available to transform crops or other host cells they may be directly applied. Accordingly, a wide variety of methods have been developed to

insert a DNA sequence into the genome of a host cell to obtain the transcription and/or translation of the sequence to effect phenotypic changes in the organism. Thus, any method that provides for efficient transformation/transfection may be employed.

5 A DNA sequence coding for the desired polynucleotide of the present invention, for example a cDNA, RNA or a genomic sequence, will be used to construct an expression cassette that can be introduced into the desired plant. Isolated nucleic acid acids of the present invention can be introduced into plants according techniques known in the art. Generally, expression cassettes as  
10 described above and suitable for transformation of plant cells are prepared.

Techniques for transforming a wide variety of higher plant species are well known and described in the technical, scientific, and patent literature. See, for example, Weising *et al.*, *Ann. Rev. Genet.* 22: 421-477 (1988). For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell  
15 using techniques such as electroporation, PEG-mediated transfection, particle bombardment, silicon fiber delivery, or microinjection of plant cell protoplasts or embryogenic callus. See, e.g., Tomes, *et al.*, Direct DNA Transfer into Intact Plant Cells Via Microprojectile Bombardment. pp.197-213 in Plant Cell, Tissue and Organ Culture, Fundamental Methods. eds. O. L. Gamborg and G.C.  
20 Phillips. Springer-Verlag Berlin Heidelberg New York, 1995. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is  
25 infected by the bacteria. See, U.S. Patent No. 5,591,616.

The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, *Embo J.* 3: 2717-2722 (1984). Electroporation techniques are described in Fromm *et al.*, *Proc. Natl. Acad. Sci.* 82: 5824 (1985). Ballistic transformation techniques are described in Klein *et al.*, *Nature* 327: 70-  
30 73 (1987).

*Agrobacterium tumefaciens*-mediated transformation techniques are well described in the scientific literature. See, for example Horsch *et al.*, *Science* 233:

496-498 (1984), and Fraley *et al.*, *Proc. Natl. Acad. Sci.* 80: 4803 (1983). For instance, *Agrobacterium* transformation of maize is described in U.S. Patent Nos. 5,550,318 and WO98/32326

Other methods of transfection or transformation include (1) *Agrobacterium*  
5 *rhizogenes*-mediated transformation (see, e.g., Lichtenstein and Fuller In: Genetic Engineering, vol. 6, PWJ Rigby, Ed., London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: DNA Cloning, Vol. II, D. M. Glover, Ed., Oxford, IRI Press, 1985), Application PCT/US87/02512 (WO 88/02405 published Apr. 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid  
10 along with *A. tumefaciens* vectors pARC8 or pARC16 (2) liposome-mediated DNA uptake (see, e.g., Freeman *et al.*, *Plant Cell Physiol.* 25: 1353, 1984), (3) the vortexing method (see, e.g., Kindle, *Proc. Natl. Acad. Sci.*, USA 87: 1228, (1990).

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, *Methods in Enzymology*, 101:433 (1983); D. Hess,  
15 *Intern Rev. Cytol.*, 107:367 (1987); Luo *et al.*, *Plane Mol. Biol. Reporter*, 6:165 (1988). Expression of polypeptide coding nucleic acids can be obtained by injection of the DNA into reproductive organs of a plant as described by Pena *et al.*, *Nature*, 325.:274 (1987). DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as described by  
20 Neuhaus *et al.*, *Theor. Appl. Genet.*, 75:30 (1987); and Benbrook *et al.*, in *Proceedings Bio Expo 1986*, Butterworth, Stoneham, Mass., pp. 27-54 (1986).

Animal and lower eukaryotic (e.g., yeast) host cells are competent or rendered competent for transfection by various means. There are several well-known methods of introducing DNA into animal cells. These include: calcium  
25 phosphate precipitation, fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of the recipient cells with liposomes containing the DNA, DEAE dextran, electroporation, biolistics, and micro-injection of the DNA directly into the cells. The transfected cells are cultured by means well known in the art. Kuchler, R.J., *Biochemical Methods in Cell Culture and Virology*, Dowden,  
30 Hutchinson and Ross, Inc. (1977).

### Transgenic Plant Regeneration

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype. Such regeneration techniques often rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with a polynucleotide of the present invention. For transformation and regeneration of maize see, Gordon-Kamm *et al.*, *The Plant Cell*, 2:603-618 (1990).

Plants cells transformed with a plant expression vector can be regenerated, e.g., from single cells, callus tissue or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant. Plant regeneration from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, Macmillan Publishing Company, New York, pp. 124-176 (1983); and Binding, *Regeneration of Plants, Plant Protoplasts*, CRC Press, Boca Raton, pp. 21-73 (1985).

The regeneration of plants containing the foreign gene introduced by *Agrobacterium* can be achieved as described by Horsch *et al.*, *Science*, 227:1229-1231 (1985) and Fraley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). This procedure typically produces shoots within two to four weeks and these transformant shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Transgenic plants of the present invention may be fertile or sterile.

Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38: 467-486 (1987). The regeneration of plants from either single plant protoplasts or various explants is well known in the art. See, for example, Methods for Plant Molecular Biology, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, Calif. (1988). For maize cell culture and regeneration see generally, *The Maize Handbook*, Freeling and Walbot, Eds.,

Springer, New York (1994); *Corn and Corn Improvement*, 3<sup>rd</sup> edition, Sprague and Dudley Eds., American Society of Agronomy, Madison, Wisconsin (1988).

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

In vegetatively propagated crops, mature transgenic plants can be propagated by the taking of cuttings or by tissue culture techniques to produce multiple identical plants. Selection of desirable transgenics is made and new varieties are obtained and propagated vegetatively for commercial use. In seed propagated crops, mature transgenic plants can be self crossed to produce a homozygous inbred plant. The inbred plant produces seed containing the newly introduced heterologous nucleic acid. These seeds can be grown to produce plants that would produce the selected phenotype.

Parts obtained from the regenerated plant, such as flowers, seeds, leaves, stems, stalks, branches, fruit, and the like are included in the invention, provided that these parts comprise cells comprising the isolated nucleic acid of the present invention. Progeny and variants, and mutants of the regenerated plants are also included within the scope of the invention, provided that these parts comprise the introduced nucleic acid sequences.

Transgenic plants expressing a selectable marker can be screened for transmission of the nucleic acid of the present invention by, for example, standard immunoblot and DNA detection techniques. Transgenic lines are also typically evaluated on levels of expression of the heterologous nucleic acid. Expression at the RNA level can be determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis can be employed and include PCR amplification assays using oligonucleotide primers designed to amplify only the heterologous RNA templates and solution hybridization assays using heterologous nucleic acid-specific probes. The RNA-positive plants can then analyzed for protein expression by Western immunoblot analysis using the specifically reactive antibodies of the present invention. In addition, *in situ* hybridization and immunocytochemistry according to standard protocols can be

done using heterologous nucleic acid specific polynucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue. Generally, a number of transgenic lines are usually screened for the incorporated nucleic acid to identify and select plants with the most appropriate expression profiles.

A preferred embodiment is a transgenic plant that is homozygous for the added heterologous nucleic acid; i.e., a transgenic plant that contains two added nucleic acid sequences, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) a heterozygous transgenic plant that contains a single added heterologous nucleic acid, germinating some of the seed produced and analyzing the resulting plants produced for altered expression of a polynucleotide of the present invention relative to a control plant (i.e., native, non-transgenic). Back-crossing to a parental plant and out-crossing with a non- transgenic plant are also contemplated.

Genotyping provides a means of distinguishing homologs of a chromosome pair and can be used to differentiate segregants in a plant population. Molecular marker methods can be used for phylogenetic studies, characterizing genetic relationships among crop varieties, identifying crosses or somatic hybrids, localizing chromosomal segments affecting monogenic traits, map based cloning, and the study of quantitative inheritance. See, e.g., *Plant Molecular Biology: A Laboratory Manual*, Chapter 7, Clark, Ed., Springer-Verlag, Berlin (1997). For molecular marker methods, see generally, *The DNA Revolution* by Andrew H. Paterson 1996 (Chapter 2) in: *Genome Mapping in Plants* (ed. Andrew H. Paterson) by Academic Press/R. G. Landis Company, Austin, Texas, pp.7-21.

The particular method of genotyping in the present invention may employ any number of molecular marker analytic techniques such as, but not limited to, restriction fragment length polymorphisms (RFLPs). RFLPs are the product of allelic differences between DNA restriction fragments caused by nucleotide sequence variability. Thus, the present invention further provides a means to follow segregation of a gene or nucleic acid of the present invention as well as

chromosomal sequences genetically linked to these genes or nucleic acids using such techniques as RFLP analysis.

Plants that can be used in the method of the invention include monocotyledonous and dicotyledonous plants. Preferred plants include corn,  
5 soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, lupin and millet.

Seeds derived from plants regenerated from transformed plant cells, plant parts or plant tissues, or progeny derived from the regenerated transformed plants, may be used directly as feed or food, or further processing may occur.

# 10 **Antibodies**

The proteins encoded by polynucleotides of this embodiment, when presented as an immunogen, elicit the production of polyclonal antibodies which specifically bind to a prototype protease inhibitor polypeptide such as, but not limited to, a polypeptide encoded by the polynucleotide of (b), *supra*, or exemplary  
15 polypeptides of SEQ ID NOS: 6, 8,10,12,14,16,18 and 20. Generally, however, a protein encoded by a polynucleotide of this embodiment does not bind to antisera raised against the prototype protease inhibitor polypeptide when the antisera has been fully immunosorbed with the reference protease inhibitor polypeptide. Methods of making and assaying for antibody binding specificity/affinity are well  
20 known in the art. Exemplary immunoassay formats include ELISA, competitive immunoassays, radioimmunoassays, Western blots, indirect immunofluorescent assays and the like.

In a preferred assay method fully immunosorbed and pooled antisera which is elicited to the prototype polypeptide can be used in a competitive binding  
25 assay to test the protein. The concentration of the prototype polypeptide required to inhibit 50% of the binding of the antisera to the prototype polypeptide is determined. If the amount of the protein required to inhibit binding is less than twice the amount of the prototype protein, then the protein is said to specifically bind to the antisera elicited to the immunogen. Accordingly, the proteins embrace  
30 allelic variants, conservatively modified variants, and minor recombinant modifications to a prototype protease inhibitor polypeptide.



## CI-2 Engineering

The amino acid sequences of the wild-type CI-2 and substituted CI-2-like polypeptides are aligned in Figure 1. Numbering of amino acid positions refers to the full length wild-type CI-2 (SEQ I.D. NO. 2) unless stated otherwise. Wild type  
5 CI-2 (from barley) contains 8 lysines, one methionine, four threonines, and one tryptophan (SEQ I.D. NO. 2). A truncated form of wild type CI-2 used in the present study (SEQ I.D. NO. 4) comprises residues 19 through 83 of the full-length wild-type plus a start methionine. Using methods known in the art for genetic and protein engineering, barley high lysine (BHL) variants with increased  
10 levels of essential amino acids were made. Preferred barley & maize variants will have increased levels of lysine, threonine, tryptophan or methionine, or combinations thereof.

BHL1 (SEQ I.D. NO. 6) contains 14 lysines. BHL2 (SEQ I.D. NO. 8) and BHL3 (SEQ. I.D. NO. 10) each contain 15 lysines. BHL1 has lysine substitutions  
15 at wild-type (SEQ I.D. NO. 2) positions 19, 34, 41, 56, 59, 62, 67, and 73 (BHL1 positions 2, 17, 24, 39, 42, 45, 50 and 56). BHL2 contains these same substitutions plus a lysine at wild-type (SEQ I.D. NO. 2) position 65 (BHL2 position 48). BHL2 also contains alanine substitutions for wild-type residues threonine-58 and glutamate-60 (threonine-41 and glutamate-43 of BHL2). The  
20 BHL3 sequence is identical to BHL2 except that these two residues at wild type positions 58 and 60 were substituted with glycine and histidine, respectively, rather than with alanine. BHL3N (SEQ. I.D. NO. 12) contains the same substitutions as BHL3, plus four lysine substitutions in the 18 additional amino acid residues in the amino terminal region, for a total of 20 lysines. The BHL4  
25 sequence (SEQ I.D. NO. 14) is the same as BHL1 except that the residue at wild type position 59 (BHL4 position 42) is glycine, rather than lysine. BHL5, BHL6, and BHL8 were designed to have an increased content of methionine, threonine, and tryptophan, as well as lysine. BHL5 (SEQ I.D. NO. 16) contains lysine substitutions at wild type positions 19, 34, 41, 47, 56, 62, 67, 73, 75, 78, and 81  
30 (BHL5 positions 3, 18, 25, 31, 40, 46, 51, 57, 59, 62, and 65). BHL5 also contains methionine substitutions at wild-type positions 17 (start methionine for BHL5), 20, 38, 40, 49, and 63, corresponding to BHL5 positions 1, 4, 22, 24, 33, and 47.

BHL5 also contains tryptophan substitutions at wild-type positions 61 and 69 (BHL5 positions 45 and 53), as well as threonine substitutions at wild-type positions 23, 31, and 79 (BHL5 positions 7, 15, and 63). BHL5 contains 17 lysines, six methionines, three tryptophans, and six threonines. BHL5 also contains the glycine substitution at wild-type position 59 (BHL5 position 43). BHL6 (SEQ. I.D. NO. 18) has the same sequence as that of BHL5, except that the residue at wild-type position 67 (BHL6 position 49) is arginine, rather than lysine. BHL8 (SEQ. I.D. NO. 20) has the same sequence as BHL6 except that cysteines were substituted at wild-type positions 22 and 82 (BHL8 positions 6 and 66).

The active site loop region encompasses an extended loop region from about amino acid residue 53 to about amino acid residue 70. Destabilization of the reactive loop was achieved by substituting the non-wild type amino acids residues at about positions 53 to about 70. Preferably, the following mutations are made (all numbering corresponds to SEQ. I.D. No. 2 unless otherwise stated): Arg62 → Lys62, Arg65 → Lys65, Arg67 → Lys67, Thr58 → Ala58 or Gly58, and Glu60 → Ala60 or His60. As an alternative approach to decreasing inhibitory activity without substantial destabilization of the active site loop, methionine 59 was changed to glycine. A glycine at this position is not known in any naturally occurring CI-2 homologs.

The first 18 residues in the wild type CI-2 do not assume any ordered conformation and also do not contribute to the structural integrity of the molecule (see e.g. Kjaer, et al., Carlsberg Res. Commun.; Vol. 53; pp. 327-354; (1987); incorporated herein in its entirety by reference), a full length 83 residue version was created in which residues one or more 1,8,11, and 17 were also replaced with one or more non-native amino acids. In one embodiment residues 1,8,11, and 17 were cysteine and conservative substitutions. In a preferred embodiment the non-native residues are methionine and lysine replaced with essential amino acids. The resulting compound has the sequence indicated in SEQ ID No. 12. Additionally, substitution of residues threonine, at position 58, and glutamic acid, at position 60, with glycine and histidine, respectively, resulted in a protein with lowered protease inhibitor activity. The resulting compound has the sequence

indicated in Sequence I.D. No. 5. The full length engineered CI-2 containing 21 lysine residues (25.3%) has also been expressed in and purified from E. coli.

In one embodiment, the CI-2-like protein has elevated essential amino acid content. Optionally, the CI-2-like protein has both elevated essential amino acid content and reduced protease inhibitor activity.

Criteria in determining sequences with homology to the present invention include determination of homology through sequence alignment using amino acids 24W, 35A, and 66V, for example and/or the amino acids 24-29, 54-58, 65-71 and/or 80-83. Alignment of these conserved residues provide a method for aligning sequences and corresponding them and their residue numbers to Seq. I.D. No. 2. Once aligned, native amino acid residues can be substituted with essential amino acids at the same residues identified as substitutable in Seq. ID No. 2.

These polypeptide comprise substituted CI-2-like polypeptides, or truncated versions thereof substituted to contain 7 or more non-native essential amino acid residues at positions corresponding to positions in Sequence ID. No.2 selected from residues 1, 8, 11, 17, 18, 19, 20, 22, 23, 31, 34, 38, 40, 41, 47, 49, 56, 58, 59, 60, 61, 62, 63, 65, 67, 69, 73, 75, 76, 78, 79, 81, 82, or combinations thereof. In another embodiment the substituted CI-2-like protein has addition non-native residues at positions 32, 45, 53, 64, 70, 74, and 77. In one embodiment the substituted CI-2-like protein has 7 or more substitutions. In another embodiment the substituted CI-2-like protein has more than 8 or more than 9 substitutions. In still another embodiment the substituted CI-2-like protein has more than 10 or more than 11. In still another embodiment the substituted CI-2-like protein has more than 14 or more than 16. In still another embodiment the substituted CI-2-like protein has more than 20 or more than 25. In still another embodiment the substituted CI-2-like protein has more than 27 or more than 30. In still another embodiment the substituted CI-2-like protein has more than 32 or more than 34. In still another embodiment the substituted CI-2-like protein has more than 35 or more than 40. In still another embodiment the substituted CI-2-like protein has more than 42 or more than 45.

In another embodiment this invention comprises a substituted CI-2-like protein with an non-native essential amino acid residue in more than about 11% to less than about 75% of the amino acid residues.

For example in figure 2 sequence 1 is aligned with CI-2 and these CI-2-like polypeptides could be substituted to contain, G19K, I38M, or R41K in accordance with the present invention. These modifications can be made using methods known in the art with the material and methods described in the instant specification.

Genes that have the desired effect are selected using procedures described in the instant specification.

In one embodiment the substituted CI-2-like protein has a non-native essential amino acid in more than about 11% to less than about 80% of the amino acid residues. In another embodiment a non-native essential amino acid residue is in more than about 12% to less than about 75% of the amino acid residues. In another embodiment a non-native essential amino acid residue is in more than about 15% to less than about 75% of the amino acid residues. In another embodiment a non-native essential amino acid residue is in more than about 15% to less than 70%. In another embodiment a non-native essential amino acid residue is in more than about 20% to less than 70%. In another embodiment a non-native essential amino acid residue is in more than about 25% to less than 65%. In another embodiment a non-native essential amino acid residue is in more than about 30% to less than 60%. In another embodiment a non-native essential amino acid residue is in more than about 50% to less than about 80% of the amino acid residues.

A substituted CI-2-like polypeptide may have from about 55 to about 90% total essential amino acid content. In one embodiment the substituted CI-2-like polypeptide has from about 60 to about 90% total essential amino acid content. In another embodiment the substituted CI-2-like polypeptide has from about 60 to about 85% total essential amino acid content. In another embodiment the substituted CI-2-like polypeptide has from about 70 to about 90% total essential amino acid content. In another embodiment the substituted CI-2-like polypeptide has 75-90% total essential amino acid content.

In one embodiment the substituted CI-2-like protein may have other modifications. In one embodiment the substituted protein has a free energy of unfolding of more than about 3.5 to about 15 Kcal/mol. In another embodiment the free energy of unfolding is more than about 4 to about 10 Kcal/mol. In another embodiment the free energy of unfolding is more than about 6 to about 10 Kcal/mol.

The substituted CI-2-like protein is made more stable by the addition of disulfide bonds. In one embodiment from one to less than 5 disulfide bonds are added. In another embodiment from one to less than 3 disulfide bonds are added. In another embodiment one disulfide bond is added. In one embodiment the disulfide bonds comprise residues [E23C and R81C] or [T22C and V82C] or [V53C and V70C]. In a preferred embodiment the disulfide bond comprises residues T22C and V82C. In another preferred embodiment the disulfide bond comprises residues E23C and R81C.

The present invention also includes the substituted CI-2-like protein with an amino terminal extension. In one embodiment the extension is for nutritional enhancement. In another embodiment the extension is a start signal, a transit sequence, a transit peptide, a signal peptide, a fusion protein, a cleavable peptide, a CI-2-like polypeptide or an uncleaved peptide. In one embodiment the CI-2 polypeptide has at least 1 to about 18 residues. In another embodiment the extension contains a nutritionally-enhancing polypeptide. In another embodiment the nutritionally-enhancing polypeptide contains essential amino acids.

The substituted CI-2-like protein with essential amino acid substitutions may also have a modified protease activity. In one embodiment the protease activity is changed in specificity.

In one embodiment of the present invention, the substituted CI-2-like protein is digestible. In one embodiment the protein is digested in simulated gastric fluid. In another embodiment the protein is digested in simulated intestinal fluid.

SEQ ID NO. 2. In another embodiment, substitutions are at 7 or more residues. In another embodiment the substituted CI-2-like protein has more than 8 or more than 9 substitutions. In still another embodiment the substituted CI-2-like protein has more than 10 or more than 11. In still another embodiment the substituted CI-2-like protein has more than 14 or more than 16. In still another embodiment the substituted CI-2-like protein has more than 20 or more than 25. In still another embodiment the substituted CI-2-like protein has more than 27 or more than 30. In still another embodiment the substituted CI-2-like protein has more than 32 or more than 34. In still another embodiment the substituted CI-2-like protein has more than 35 or more than 40. In still another embodiment the substituted CI-2-like protein has more than 42 or more than 45.

In one embodiment the substituted CI-2-like protein exhibits reduced inhibiting activity against chymotrypsin, subtilisin and elastase. In another embodiment the substituted CI-2-like protein exhibits no inhibitory activity against chymotrypsin and elastase.

In one embodiment the substituted CI-2-like protein has 2 or more or 3 or more substitutions. In another embodiment the substituted CI-2-like protein has more than 4 or more than 5 substitutions. In still another embodiment the substituted CI-2-like protein has more than 7 or more than 9. In still another embodiment the substituted CI-2-like protein has more than 10 or more than 11. In still another embodiment the substituted CI-2-like protein has more than 12 or more than 15. In still another embodiment the substituted CI-2-like protein has more than 17 or more than 20. In still another embodiment the substituted CI-2-like protein has more than 22 or more than 24. In still another embodiment the substituted CI-2-like protein has more than 25 or more than 27. In still another embodiment the substituted CI-2-like protein has more than 30 or more than 35.

In one embodiment an essential amino acid is methionine, threonine, lysine, isoleucine, leucine, valine, tryptophan, phenylalanine, and histidine. In another embodiment the essential amino acid is lysine, threonine, tryptophan, methionine, or combinations and conservative substitutions thereof.

The following conservative essential amino acid substitutions are included in the present invention: [M, I, L, V] or [K, T]. K is replaceable with T. M, I, L and V are replaceable with each other.

For example selection of [E34K] and [I56M, T58G, M59G, E60H, Y61W, R62K] provides substituted CI-2-like polypeptide having the residues of SEQ ID NO. 2 at all positions except 34, 56, 58, 59, 60, 61 and 62 where amino acids are K, M, G, G, H, W, & K, respectively.

Nutritional enhancement may also be provided through insertion into the active site loop region. In one embodiment this insert is one or more of a combination of essential amino acids.

In a preferred embodiment the insert is a peptide of from 2 to 20 amino acids. In another embodiment the peptide is from 5 to 15 amino acids. In another embodiment the essential amino acids are lysine, threonine, methionine or tryptophan or combinations thereof.

One embodiment of the present invention provides an isolated polypeptide comprising a plant substituted CI-2-like polypeptide having the following composition: 15-35 mole % lysine, 5-15 mole % methionine, 6-25 mole % threonine, 4-9 mole % tryptophan or combinations thereof. In another embodiment the plant substituted CI-2-like polypeptide has the following composition: 20-35 mole % lysine, 7-15 mole % methionine, 10-25 mole % threonine, 6-9 mole % tryptophan or combinations thereof.

In one embodiment the substituted CI-2-like polypeptide is proteolytically stable, as demonstrated by detection of the intact polypeptide based upon detection on SDS-PAGE gel, following a 30 minute incubation at 37°C in 100mM Tris-HCl, 50mMNaCl, 1mM $\text{CaCl}_2$ , pH 8, with a 10:1 (weight to weight ratio) of polypeptide:protease, with the protease being either chymotrypsin or trypsin.

In one embodiment of the present invention an isolated polypeptide comprises at least 23 contiguous amino acids with more than 79% sequence identity, to the polypeptide of Seq. ID No. 20, wherein the % sequence identity is based on the 23 contiguous amino acids sequence and is determined by GAP analysis using Gap Weight of 8 and Length Weight of 2. In another embodiment an isolated polypeptide comprises at least 23 contiguous amino acids with more

than 81% sequence identity, to the polypeptide of Seq. ID No. 20. In another embodiment an isolated polypeptide comprises at least 23 contiguous amino acids with more than 83% sequence identity, to the polypeptide of Seq. ID No. 20. In another embodiment an isolated polypeptide comprises at least 23 contiguous amino acids with more than 85% sequence identity, to the polypeptide of Seq. ID No. 20. In another embodiment an isolated polypeptide comprises at least 23 contiguous amino acids with more than 89% sequence identity, to the polypeptide of Seq. ID No. 20.

In one embodiment of the present invention, the polynucleotide has at least 73% sequence identity to SEQ ID NO: 19, wherein the % sequence identity is based on the entire sequence and is determined by BLAST 2.0. In another embodiment the polynucleotide has at least 75% or 77% sequence identity to SEQ ID NO: 19. In another embodiment the polynucleotide has at least 80% or 85% sequence identity to SEQ ID NO: 19. In another embodiment the polynucleotide has at least 90% or 95% sequence identity to SEQ ID NO: 19. In another embodiment the polynucleotide has 98 sequence identity to SEQ ID NO: 19.

In an embodiment of the present invention, the polynucleotide comprising at least 25 nucleotides in length which hybridizes under low stringency conditions to a polynucleotide having the sequence set forth in SEQ ID NOs: 19, wherein the conditions include hybridization with a buffer solution of 30% formamide, 1 M NaCl, 1% SDS at 37°C for 24 hours and a wash in 2X SSC at 50°C, 3x for 15 minutes.

Modification in the active site loop area by amino acid substitution or other means, destroys the hydrogen bonding and changes or reduces the protease inhibitor activity of BHL. Substitution of amino acid residues threonine, at position 58, and glutamic acid, at position 60, with glycine and histidine, respectively, resulted in a protein with lowered protease inhibitor activity. Residue 59, when changed, is able to modifying protease inhibitor activity and change specificity. When this residue was changed to a lysine, the protease inhibition specificity was changed from a chymotrypsin inhibitor to a trypsin inhibitor. When residue 59 was changed to glycine, the inhibitory activity against trypsin was removed, and



inhibitory activity against chymotrypsin, subtilisin, and elastase was considerably reduced compared to wild type CI-2.

## 5 **Proteins**

Synthesis of the compounds is performed according to methods of peptide synthesis which are well known in the art and thus constitute no part of this invention. For example, in vitro, the compounds can be synthesized on an applied Biosystems model 431a peptide synthesizer using fastmoc™ chemistry involving hbtu [2-(1h-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, as published by Rao, et al., Int. J. Pep. Prot. Res.; Vol. 40; pp. 508-515; (1992); incorporated herein in its entirety by reference. Peptides can be cleaved following standard protocols and purified by reverse phase chromatography using standard methods. The amino acid sequence of each peptide can be confirmed by automated edman degradation on an applied biosystems 477a protein sequencer/120a pth analyzer. More preferably, however, the compounds of this invention are synthesized in vivo by bacterial or plant cells which have been transformed by insertion of an expression cassette containing a synthetic gene which when transcribed and translated yields the desired compound. Such empty expression cassettes, providing appropriate regulatory sequences for plant or bacterial expression of the desired sequence, are also well-known, and the nucleotide sequence for the synthetic gene, either RNA or DNA, can readily be derived from the amino acid sequence for the protein using standard reference texts. Preferably, such synthetic genes will employ plant-preferred codons to enhance expression of the desired protein.

Promoters that may be used in the genetic sequence include NOS, OCS and CaMV promoters.

This invention provides a method for increasing essential amino acid levels in Agrobacterium tumefaciens-susceptible dicotyledonous plants in which the expression cassette is introduced into the cells by infecting the cells with Agrobacterium tumefaciens, a plasmid of which has been modified to include a

plant expression cassette of this invention. Agrobacterium tumefaciens-mediated transformation is also effective for monocotyledonous plants.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention  
5 pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Variations on the above embodiments are within the ability of one of  
10 ordinary skill in the art, and such variations do not depart from the scope of the present invention as described in the following claims.

The present invention will be further described by reference to the following detailed examples. It is understood, however, that there are many extensions, variations, and modifications on the basic theme of the present  
15 invention beyond that shown in the examples and description, which are within the spirit and scope of the present invention. All publications, patents, and patent applications cited herein are hereby incorporated by reference.

#### **Assays for Compounds that Modulate Protease Inhibitory Activity or Expression**

20 The present invention also provides means for identifying compounds that bind to (e.g., substrates), and/or increase or decrease (i.e., modulate) the inhibitory activity of, protease inhibitor polypeptides. The method comprises contacting a protease inhibitor polypeptide of the present invention with a compound whose ability to bind to or modulate inhibitory activity is to be  
25 determined. The protease inhibitor polypeptide employed will have at least 20%, preferably at least 30% or 40%, more preferably at least 50% or 60%, and most preferably at least 70% or 80% of the inhibitory activity of the full-length (native and endogenous) protease inhibitor polypeptide. Generally, the protease inhibitor polypeptide will be present in a range sufficient to determine the effect of  
30 the compound, typically about 1 nM to 10  $\mu$ M. Likewise, the compound will be present in a concentration of from about 1 nM to 10  $\mu$ M. Those of skill will understand that such factors as enzyme concentration, ligand concentrations

6650" 69TCE50

(i.e., substrates, products, inhibitors, activators), pH, ionic strength, and temperature will be controlled so as to obtain useful kinetic data and determine the presence or absence of a compound that binds or modulates protease inhibitor polypeptide activity. Methods of measuring enzyme kinetics is well known in the art. See, e.g., Segel, *Biochemical Calculations*, 2<sup>nd</sup> ed., John Wiley and Sons, New York (1976).

Although the present invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

### **Examples**

#### **Example 1- Construction of Expression Cassettes**

Vector construction was based upon the published WT CI-2A sequence information Williamson *et al*, Eur. J. Biochem 165: 99-106 (1987) and SEQ ID NO 1. Methods for obtaining full length or truncated wild-type CI-2 DNA include, but are not limited to PCR amplification, from a barley (or other plant) endosperm cDNA library using oligonucleotides derived from Seq. ID No 1 or from the published sequence *supra*, using probes derived from the same on a barley endosperm cDNA library, or using a set of overlapping oligonucleotides that encompass the gene, or having the gene synthesized by a commercial vendor such as The Midland Certified Reagent Company (Midland, Texas).

#### **BHL1**

The BHL1 insert corresponds to SEQ ID NO 5. Oligonucleotide pairs, N4394/N4395, and N4396/N4397, were annealed and ligated together to make a 202 base pair double stranded DNA molecule with overhangs compatible with *Rca* I and *Nhe* I restriction sites. PCR was performed on the annealed molecule using primers N5045 and N5046 to add a 5' *Spe* I site and 3' *Hind* III site. The PCR product was then restriction digested at those sites and ligated into pBluescript II KS+ at *Spe* I and *Hind* III sites. The insert was then removed by

restriction digestion with *Rca* I and *Hind* III and was ligated into the *Nco* I and *Hind* III sites of pET28a (Novagen) to form the BHL1 construct.

Oligonucleotide sequences (5' to 3'):

N4394

5        1     CATGAAGCTG   AAGACAGAGT   GGCCGGAGTT   GGTGGGGAAA  
TCGGTGGAGA

      51     AAGCCAAGAA   GGTGATCCTG   AAGGACAAGC   CAGAGGCGCA  
AATCATAGTT

     101 CTGC

10     N4395

      1     CAACCGGCAG   AACTATGATT   TGCGCCTCTG   GCTTGTCTT  
CAGGATCACC

     51     TTCTTGGCTT   TCTCCACCGA   TTTCCCCACC   AACTCCGGCC  
ACTCTGTCTT

15        101 CAGCTT

      N4396

      1     CGGTTGGTAC   AAAGGTGACG   AAGGAATATA   AGATCGACCG  
CGTCAAGCTC

20        51 TTTGTGGATA AAAAGGACAA CATCGCGCAG GTCCCCAGGG TCGG

      N4397

      1     CTAGCCGACC   CTGGGGACCT   GCGCGATGTT   GTCCTTTTTA  
TCCACAAAGA

25        51 GCTTGACGCG GTCGATCTTA TATTCCTTCG TCACCTTTGT AC

      N5045

      1     GTACTAGTCA TGAAGCTGAA GACAGA

30     N5046

      1     GAGAAGCTTG CTAGCCGACC CTGGGGAC

## BHL2

The BHL2 construct insert corresponds to SEQ ID NO 7. An overlap PCR strategy was used to make the BHL2 construct. PWO polymerase from Boehringer-Mannheim was used for all PCR reactions. The primers were chosen to change 3 amino acids in the BHL1 active site loop region, and to create unique *Age* I and *Hind* III restriction sites flanking the active site loop, to facilitate loop replacement in future constructs. A unique *Rca* I site (compatible with *Nco* I) was included at the 5' end, and a unique *Xho* I site was included at the 3' end. The overlap PCR was done as follows: PCR was done with primers N13561 and N13564, using the BHL1 construct as template. A separate PCR was done with primers N13563 and N13562, again using the BHL1 construct as template. The products from both reactions were gel purified and combined. Primer N13565, which overlapped regions on both of the PCR products, was then added and another PCR was done to generate the full-length insert. The resulting product was amplified by another PCR with primers N13561 and N13562. It was subsequently suspected that a deletion was present in N13562 that caused a frameshift near the 3' end of the PCR product. To avoid this frameshift problem, a final PCR reaction was done with primers N13562 and N13905. The final PCR product was digested with *Rca* I and *Xho* I, and then ligated into the *Nco* I and *Xho* I sites of pET 28b. Note: Some primers had 6-oligonucleotide extensions to improve restriction digestion efficiency.

Oligonucleotide sequences (5' TO 3'):

N13561

1 TTTTTTTCATGAAGCTGAAGACA

N13562 (as ordered)

1 TTTTTTCTCGAGGCTAGCCGACCCTGGGGA

N13563

1 ATCGACAAGGTCAAGCTTTTTGTGGATAAAAAGGA

N13564

1 CACCTTTGTACCAACCGGTAGAACTATGATTTGCGC

N13565

1 GTTGGTACAAAGGTGGCGAAGGCCTATAAGATCGACAAGGTCAAG

N13905

5 1 TTTTTTCTCGAGGCTAGCCGACCCTGGGGACCTGCGCTA

### **BHL3**

The BHL3 construct insert corresponds to SEQ ID NO 9. The BHL2 construct was digested with *Age* I and *Hind* III, and the region between these sites was removed by gel purification and discarded. Oligonucleotide pairs, N14471 and N14472, were annealed to make a double stranded DNA molecule with overhangs compatible with *Age* I and *Hind* III restriction sites. The annealed product was ligated into the *Age* I and *Hind* III sites of the digested BHL2 construct to yield the BHL3 construct.

15 Oligonucleotide sequences (5' to 3'):

N14471

1 CCGGTTGGTACAAAGGTGGGTAAGCATTATAAGATCGACAAGGTCA

N14472

20 1 AGCTTGACCTTGTCGATCTTATAATGCTTACCCACCTTTGTACCAA

### **BHL3N**

The BHL3N construct insert corresponds to SEQ ID No 11. A PCR reaction was done with the BHL3 construct as template. The primers for this reaction were N13771 and N13905. The resulting PCR product was digested with *Rca* I and *Xho* I and ligated into the *Nco* I and *Xho* I sites of pET 28b to yield the BHL3N construct.

Oligonucleotide sequences (5' to 3'):

N13771

30 1  
TTTTTTTCATGAAGTCGGTGGAGAAGAAACCGAAGGGTGTGAAGACAGGTGC  
GGGTGACAAGCATAAGCTGAAGACAGAGTG

N13905 (already provided in BHL2 description).

#### **BHL4**

The BHL4 construct insert DNA corresponds to SEQ ID NO 13. The BHL2 construct was digested with *Age* I and *Hind* III, and the region between these sites was removed by gel purification and discarded. Oligonucleotide pairs, N22098 and N22099, were annealed to make a double stranded DNA molecule with overhangs compatible with *Age* I and *Hind* III restriction sites. The annealed product was ligated into the *Age* I and *Hind* III sites of the digested BHL2 construct to yield the BHL4 construct.

Oligonucleotide sequences (5' to 3'):

N22098

1 CCGGTTGGTACAAAGGTGACGGGCGAATACAAGATCGACCGCGTCA

N22099

1 AGCTTGACGCGGTTCGATCTTGTATTGCCCCGTACCTTTGTACCAA

#### **BHL5**

The BHL5 construct insert DNA corresponds to SEQ ID NO 15. This gene was synthesized by a commercial vendor, The Midland Certified Reagent Company (Midland, Texas). The gene was supplied by Midland following digestion by *Nco* I and *Hind* III, and was ligated into the *Nco* I and *Hind* III sites of pET 28b to yield the BHL5 construct.

#### **BHL6**

The BHL6 construct insert DNA corresponds to SEQ ID NO 17. The BHL5 construct was digested with *Age* I and *Sa* I, and the region between these sites was removed by gel purification and discarded. Oligonucleotide pairs, N23923 and N23924, were annealed to make a double stranded DNA molecule with overhangs compatible with *Age* I and *Sa* I restriction sites. The annealed product was ligated into the *Age* I and *Sa* I sites of the digested BHL5 construct to yield the BHL6 construct.

Oligonucleotide sequences (5' to 3'):

N23923

1 CCGGTGAATGGAAGATGGATCGCGTCCGCCTCTGGG

N23924

1 TCGACCCAGAGGCGGACGCGATCCATCTTCCATTCA

5

### **BHL8**

The BHL8 construct insert DNA corresponds to SEQ ID No 19. A PCR reaction was done using the BHL6 construct as template. The primers for this reaction were N26671 and N26672. The resulting PCR product was digested with *Nco* I and *Hind* III and ligated into the *Nco* I and *Hind* III sites of pET 28b to yield the BHL8 construct.

Oligonucleotide sequences (5' to 3'):

N26671

1 TTTTTCATGGCTAAGATGAAGTGCACGTGGCCTGAGCTGGT

N26672

1 TTTTTTAAGCTTGGATCCCTAGCCGCACTTCGGAGTCTTGGCGA

The following experiments used truncated wild type CI-2.

### **Example 2 - Expression of BHL Proteins in *E. coli*, Purification, and Verification of Recombinant Protein Sequence**

#### ***Expression in E. coli***

BHL1, BHL2, BHL3, BHL3N, BHL4, BHL5, BHL6, BHL8, and the truncated wild-type CI-2 were expressed in *E. coli* using materials and methods from Novagen, Inc. The Novagen expression vector pET-28 was used (pET-28a for WT CI-2 and BHL1, and pET-28b for the other proteins). *E. coli* strains BL21(DE-3) or BL21(DE-3)pLysS were used. Cultures were typically grown until an OD at 600 nm of 0.8 to 1.0, and then induced with 1 mM IPTG and grown another 2.5 to 5 hours before harvesting. Induction at an OD as low as 0.4 was also done successfully. Growth temperatures of 37 degrees centigrade and 30 degrees centigrade were both used successfully. The media used was 2xYT plus the appropriate antibiotic at the concentration recommended in the Novagen manual.



### *Purification*

a. WT Cl-2 (truncated)-- Lysis buffer was 50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 150 mM NaCl. The protein was precipitated with 70% ammonium sulfate. The pellet was dissolved and dialyzed against 50 mM Tris-HCl, pH 8.6. The protein was loaded onto a Hi-Trap Q column, and the unbound fraction was collected and precipitated in 70% ammonium sulfate. The pellet was dissolved in 50 mM sodium phosphate, pH 7.0, 200 mM NaCl, and fractionated on a Superdex-75 26/60 gel filtration column. Fractions were pooled and concentrated.

b. BHL1--Lysis buffer was 50 mM sodium phosphate, pH 7.0, 1 mM EDTA. The protein was loaded onto an SP Sepharose FF 16/10 column, washed with 150 mM NaCl in 50 mM sodium phosphate, pH 7.0, and then eluted with an NaCl gradient in 50 mM sodium phosphate. BHL1 eluted at approximately 200 mM NaCl. Fractions were pooled and concentrated.

c. BHL2, BHL3, BHL3N, BHL4, BHL5, BHL6, and BHL8--Lysis buffer was 50 mM Hepes, pH 8.0, 2mM EDTA, 0.1% Triton X-100, and 0.5 mg/ml lysozyme. The protein was loaded onto an SP-Sepharose cation exchange column (typically a 5 to 10 ml size), washed with 50 mM sodium phosphate, pH 7.0, and step eluted with increasing concentrations of NaCl in 50 mM sodium phosphate, pH 7.0. The protein was concentrated and then subjected to Superdex-75 gel filtration chromatography. The Superdex chromatography was done in 50 mM Tris-HCl, 150mM NaCl, pH 8.6 for BHL8, and in 50 mM sodium phosphate, 150 mM NaCl, pH 7.0 for the other proteins.

### *Storage*

The purified proteins were stored long term by freezing in liquid nitrogen and keeping frozen at -70 degrees centigrade.

### *Verification of recombinant protein sequence*

#### a. DNA sequencing--

The insert region of these pET 28 constructs was confirmed by DNA sequencing.

#### b. N-terminal protein sequencing --

100 µg of purified BHL3 were digested with 1µg of chymotrypsin (Sigma catalog # C-4129) for 30 min at 37 degrees centigrade in 50 mM sodium phosphate, pH

7.0. The resulting chymotryptic fragments were purified by reversed phase chromatography, using an acetonitrile gradient for elution. Three pure peaks were observed and were sent to the University of Michigan Medical School Protein Structure Facility for N-terminal sequencing (6 cycles). Peak 1 had an N-terminal sequence of val-asp-lys-lys-asp-asn. Peak 2 had an N-terminal sequence of lys-ile-asp-lys-val-lys. Peak 3 had an N-terminal sequence of met-lys-leu-lys-thr-glu. These results demonstrate that chymotrypsin cleaved BHL3 after tyr-61 and phe-69. The N-terminal sequences all match exactly the BHL3 expected sequence, assuming that the start methionine was largely retained in the recombinant protein. This experiment verifies that the protein we expressed in and purified from *E. coli* was BHL3.

160 µg of BHL3N were digested with 1.6 µg pepsin overnight, and the resulting peptic fragments were purified by reversed phase chromatography. Five of the resulting peaks were sent to the Iowa State University Protein Facility for N-terminal sequencing through four cycles. The N-terminal sequences of the 5 peaks were: val-gly-lys-ser, phe-val-asp-lys, pro-val-gly-thr, met-lys-ser-val, and ile-ile-val-leu, all of which exactly match the expected BHL3N sequence, assuming that the start methionine was retained in this recombinant protein. This experiment verifies that the protein we expressed in and purified from *E. coli* was BHL3N. Samples of the other purified proteins were also subjected to N-terminal sequencing. The truncated wild type CI-2 sequence (through four cycles) was Met-Asn-Leu-Lys, as predicted from the DNA sequence. The sequence for BHL1, BHL2, and BHL4 was Met-Lys-Leu-Lys, again confirming the identity of these proteins. The sequence for BHL5, BHL6, and BHL8 was Ala-Lys-Met-Lys, again confirming the identity of these proteins but also revealing that the start methionine was not retained in these three proteins when expressed in *E. coli*.

#### c. Mass spectrometry--

All of the purified proteins were subjected to analysis by mass spectrometry. The determined masses and the predicted masses were very similar, further confirming the sequence of the engineered proteins.

### **Example 3 – Addition of Disulfide Bonds.**



again examined using this method. The free energy of unfolding in the absence of denaturant ( $\Delta G_{H_2O}$ ) and the guanidium chloride concentration sufficient for 50% unfolding are presented in the following tables.

- 5 *Equilibrium unfolding parameters (mean  $\pm$  standard deviation). Unfolding was monitored by the change in fluorescence intensity.*

Protein	$\Delta G_{H_2O}$ (kcal mol <sup>-1</sup> )	[GdmCl] <sub>50%</sub> (M)
WT CI-2	7.04 $\pm$ 0.04	3.97 $\pm$ 0.01
BHL1	4.48 $\pm$ 0.34	2.36 $\pm$ 0.04
BHL2, BHL3, & BHL3N (pooled)	1.56 $\pm$ 0.16	0.86 $\pm$ 0.02
BHL4	4.93 $\pm$ 0.19	2.59 $\pm$ 0.01

- 10 *Equilibrium unfolding parameters (mean  $\pm$  standard deviation). Unfolding was monitored by change in CD spectra at 234 nm.*

Protein	$\Delta G_{H_2O}$ (kcal mol <sup>-1</sup> )	[GdmCl] <sub>50%</sub> (M)
WT CI-2	7.52 $\pm$ 0.52	3.86 $\pm$ 0.02
BHL4	4.49 $\pm$ 0.39	2.67 $\pm$ 0.01
BHL5	2.20 $\pm$ 0.23	1.32 $\pm$ 0.05
BHL6	3.09 $\pm$ 0.08	1.78 $\pm$ 0.01
BHL8	6.96 $\pm$ 0.72	3.61 $\pm$ 0.02
BHL8 (reduced)	2.35 $\pm$ 0.10	1.66 $\pm$ 0.02

- 15 These results show that the disulfide bond of BHL8 unexpectedly led to a significantly increased thermodynamic stability of this protein over the non-disulfide bonded counterpart BHL6. When the experiment was performed with

BHL8 that had first been treated with 10 mM dithiothreitol to reduce the disulfide bond, the stability was decreased to a value less than that of BHL6. This confirmed that it was the disulfide bond of BHL8, and not just the two cysteine substitutions, that increased the thermodynamic stability of BHL8 over BHL6.

5 **Example 5 – Proteolytic Stability of Engineered Proteins, and Increased Stability Achieved by Addition of a Disulfide Bond.**

Stability of engineered proteins in the presence of proteases such as trypsin or chymotrypsin can provide insights on structural integrity of the proteins. Malformed proteins tend to be less proteolytically stable than compact, correctly folded proteins. Trypsin and chymotrypsin digests of BHL1, BHL2, BHL3, BHL3N, BHL4, and wild type CI-2 were done for 30 min at 37°C. Three micrograms of WT or engineered CI-2 were incubated with 0.3 µg protease in 100 mM Tris-HCl, 50 mM NaCl, 1mM CaCl<sub>2</sub>, pH 8.0, in a volume of 15 µl. Control samples with protease only were incubated in the same buffer. Reactions were stopped by adding an equal volume of Bio-Rad 2X Tris-Tricine SDS sample buffer containing 6 mM PMSF, followed by boiling 5 min. and then analysis by SDS-PAGE. Results are summarized in the following table:

Intact protein detectable after 30 minute incubation with trypsin or chymotrypsin.

	Trypsin	Chymotrypsin
Wild type CI-2	Yes	Yes
BHL1	Yes	Yes
BHL2	No	No
BHL3	No	No
BHL3N	No	No
BHL4	Yes	Yes

WT CI-2 and BHL1 were resistant to trypsin, and BHL4 was unexpectedly partially resistant, with some intact BHL4 protein remaining after 30 min. The other proteins were completely digested by trypsin into fragments too small to be detected by SDS-PAGE. With respect to chymotrypsin, WT CI-2 was completely resistant, as is to be expected for a chymotrypsin inhibitor. BHL1 and BHL4 were

partially resistant whereas derivatives BHL2, BHL3 and BHL3N were completely digested into smaller fragments, with no intact protein remaining.

Using the same buffer and substrate to protease ratio, BHL5, BHL6, BHL8, BHL4, and wild-type CI-2 were incubated with trypsin for 2 min, 4 min, 8 min, 15 min, 30 min, 60 min, or 120 min, or with chymotrypsin for 1 min, 2 min, 4 min, 8 min, 15 min, 30 min, 60 min, or 120 min. Results are summarized in the following table.

Longest time that intact protein still remained during incubation with trypsin or chymotrypsin.

	Trypsin	Chymotrypsin
Wild type CI-2	120 min	120 min
BHL4	60 min	120 min
BHL5	< 2 min	1 min
BHL6	2 min	4 min
BHL8	120 min	120 min
BHL8 (reduced)	< 2 min	1 min

With respect to trypsin, intact protein was still detected for BHL8 and for wild type CI-2 at 120 min., for BHL4 at 60 min, and for BHL6 at 2 min. No BHL5 was detected even at 2 min. With respect to chymotrypsin, intact protein was still detected for wild type CI-2, BHL8, and BHL4 at 120 min., for BHL6 at 4 min., and for BHL5 at 1 min. The same experiment was also done with BHL8 that had first been treated with 10 mM DTT 1 hour at 37 degrees centigrade to reduce the disulfide bond. Reduced BHL8 was not detectable even at 2 min with trypsin, and was detected only at 1 min. with chymotrypsin. This confirms that it is the disulfide bond of BHL8, and not just the cysteine substitutions, that are responsible for the increased proteolytic stability of BHL8 compared to BHL6.

In contrast to the results with BHL8, addition of the same disulfide bond in BHL3 (i.e. between cysteines substituted for Thr-22 and Val-82) did not improve the stability of BHL3 against trypsin. This experiment was done in the same buffer as described above, but with a 1:100 ratio of trypsin to substrate protein,

rather than a 1:10 ratio. BHL3 with or without the disulfide was somewhat detectable at 15 min., but not at 60 min.

The relative proteolytic stability of BHL8, BHL4, and BHL1 evident here may prove beneficial. These proteolytically stable proteins may be relatively resistant to plant proteases, which may allow them to accumulate to useful levels in plants. Furthermore, when eaten by ruminants such as cattle or sheep, proteolytically stable proteins may have a better chance of resisting digestion by bacteria in the rumen. The proteins would then be subsequently available to the animal following passage out of the rumen (McNabb et al, J. Sci. Food Agric. 64: 53-61 (1994)). The stability against trypsin and chymotrypsin does not necessarily mean that these proteins would be poorly digested by monogastric animals, because the proteins would first have to pass through the stomach, where digestion by pepsin could potentially occur, before they encounter trypsin or chymotrypsin in the intestine.

#### **Example 6 – Digestibility of Engineered Proteins in Simulated Gastric Fluid and Simulated Intestinal Fluid.**

##### *Digestion in simulated gastric fluid.*

How quickly a protein is digested in simulated gastric fluid may be an indication of how easily digestible it would be in the stomach of an animal or human. Furthermore, proteins that are quickly digested in simulated gastric fluid are less likely to be food allergens than are proteins that are stable in simulated gastric fluid (Astwood et al, Nature Biotechnology 14: 1269-1271, (1996)). Digestibility of the BHL proteins was assessed at 37 degrees centigrade in simulated gastric fluid (34 mM NaCl, 0.7% HCl, and 3.2 mg/ml pepsin). Porcine stomach pepsin (Sigma cat # P-6887) was used. Aliquots of the incubation mix containing 3 µg of wild type or engineered CI-2 in 15 to 20 µl were removed at various times and assessed by SDS-PAGE. Time points of 15 sec, 30 sec, 1 min, 5 min, and 30 min were used for wild type CI-2, BHL1, BHL2, BHL3, BHL3N, and BHL4. All of these proteins were digested in simulated gastric fluid within 15 seconds. In separate experiments, time points used for BHL5, BHL6, BHL8, BHL4 (repeat) and wild type CI-2 (repeat) were 30 sec, 1 min, 2 min, 4 min, 8

min, 15 min, and 30 min. All of these proteins were digested in simulated gastric fluid within 30 seconds. It therefore appeared that all of the BHL proteins and wild type CI-2 were easily digested by pepsin in simulated gastric fluid. In contrast to the proteins of the present study, the soybean Kunitz trypsin inhibitor  
5 was stable for one hour in simulated gastric fluid (Astwood et al, Nature Biotechnology 14: 1269-1271, (1996)).

#### *Digestion in simulated intestinal fluid.*

Simulated intestinal fluid was prepared by dissolving 68 mg of monobasic potassium phosphate in 2.5 ml of water, adding 1.9 ml of 0.2 N sodium hydroxide  
10 and 4 ml of water. Then 2.0 g porcine pancreatin (Sigma catalog # P-7545) was added and the resulting solution was adjusted with 0.2N sodium hydroxide to a pH of 7.5. Water was added to make a final volume of 10 ml. 50 µl of 1mg/ml BHL3N or wild-type CI-2 were incubated with 250 µl simulated intestinal fluid at 37 degrees centigrade . At 15 sec, 30 sec, 1 min, 5 min, and 30 min, 40 µl  
15 aliquots were removed and added to 40 µl of a stop solution consisting of 2X Tris-Tricine SDS sample buffer (Biorad) containing 2 mM EDTA and 2mM phenylmethylsulfonyl fluoride (Sigma catalog # P-7626). Digestion was assessed by 16.5 % Tris-Tricine SDS-PAGE (precast gels from Biorad). BHL3N was digested by simulated intestinal fluid within 15 seconds. In contrast, wild type CI-  
20 2 was resistant to digestion for 30 minutes. This experiment shows that in the intestine of humans or monogastric animals, the intact engineered protein would likely be more digestible than the intact wild type protein would be. Considering the previous experiments with simulated gastric fluid, however, it may be that little of either the wild type or engineered proteins would escape digestion by pepsin in  
25 the stomach to reach the intestine intact.

#### **Example 7 - Protease Inhibition Assays**

The following proteases were used to measure inhibition with CI-2 and the mutants: bovine pancreatic chymotrypsin (Sigma # C-4129), bovine pancreatic  
30 trypsin (Sigma # T-8918), porcine pancreatic elastase (Sigma # E-0258), and Subtilisin Carlsberg from *Bacillus licheniformis* (Sigma # P-5380). Assays were done at 37°C for chymotrypsin, and at 25°C for the other proteases. Reaction



volumes were typically 200  $\mu$ l and were started by addition of substrate, following preincubation for 15 min with elastase and 30 min with the other proteases. Chymotrypsin and subtilisin assays were done in 200 mM Tris-HCl, pH 8.0, with 1 nM protease and 1  $\mu$ M WT or engineered CI-2, using 1 mM N-Succinyl-Ala-Ala-Pro-Phe-p Nitroanilide (Sigma #5-7388) as substrate. Trypsin assays were done in 50 mM Tris-HCl, 2 mM NaCl, 2 mM  $\text{CaCl}_2$ , 0.005% TritonX-100, pH 7.5, with 0.5 nM trypsin and 5  $\mu$ M WT or engineered CI-2. The substrate was 1 mM N-Benzoyl-2-Ile-Glu-Gly-Arg-p-Nitroanilide (Chromogenix S-2222). Elastase assays were done in 200 mM Tris-HCl, pH 8.0 with 50 nM elastase and 2  $\mu$ M WT or engineered CI-2. The substrate was 1 mM N-succinyl-Ala-Ala-Ala-p-Nitroanilide (Sigma #S-4760). The linear increase in absorbance at 405 nm was monitored over time. Activities in the presence of WT or engineered CI-2 were expressed as a percentage of the activity of the uninhibited proteases. The results are summarized in the following table.

*Protease activity in the presence of WT or engineered CI-2. Values are expressed as a per cent of control assays containing no WT or engineered CI-2 (mean  $\pm$  standard deviation).*

Protein	Protease activity (% of control)			
	Chymotrypsin	Subtilisin	Trypsin	Elastase
WT CI-2	9 $\pm$ 4	0.3 $\pm$ 0.4	105 $\pm$ 6	3 $\pm$ 1
BHL1	87 $\pm$ 6	15 $\pm$ 2	14 $\pm$ 4	104 $\pm$ 5
BHL2	97 $\pm$ 13	82 $\pm$ 5	91 $\pm$ 8	107 $\pm$ 5
BHL3	102 $\pm$ 5	101 $\pm$ 9	104 $\pm$ 6	107 $\pm$ 7
BHL3N	98 $\pm$ 10	96 $\pm$ 2	108 $\pm$ 4	105 $\pm$ 5
BHL4	73 $\pm$ 10	50 $\pm$ 3	100 $\pm$ 4	104 $\pm$ 11
BHL5	101 $\pm$ 8	57 $\pm$ 8	101 $\pm$ 13	106 $\pm$ 0.1
BHL6	101 $\pm$ 8	37 $\pm$ 3	98 $\pm$ 2	109 $\pm$ 4
BHL8	102 $\pm$ 7	35 $\pm$ 1	111 $\pm$ 4	107 $\pm$ 2

The wild type protein was an effective inhibitor of chymotrypsin, subtilisin, and elastase, but not of trypsin, consistent with a previous study (Longstaff et al., Biochemistry 29: 7339-7347, (1990)). Compared to wild type CI-2, the engineered proteins have reduced inhibitory activity against chymotrypsin, subtilisin, and elastase. Except for BHL1, the engineered proteins also are not effective inhibitors of trypsin. A further experiment was done with BHL4. This protein was first digested with pepsin for 30 seconds, and then the inhibitory activity of the peptic fragments was assessed against chymotrypsin or subtilisin. The BHL4 peptic fragments retained no inhibitory activity against either chymotrypsin or subtilisin.

### **Example 8 - Protein Conformation**

#### **Analysis of Engineered Proteins by Circular Dichroism**

The wild-type and engineered proteins were analyzed by far UV circular dichroism (CD) spectroscopy in 10mM sodium phosphate, pH 7. The CD spectra for BHL1, BHL2, BHL3, BHL3N, and BHL4 were very similar to that of wild-type CI-2, suggesting that these proteins have similar secondary structures. The spectra for BHL5, BHL6, and BHL8 were also similar overall to the WT CI-2 spectrum, but with detectable increases in ellipticity values for BHL5 and BHL8. The wild-type protein and BHL5, BHL6, and BHL8 were also analyzed by near UV (250 nm to 350 nm) circular dichroism spectroscopy. Differences in the BHL8 spectrum were detected relative to the others.

#### **Example 9 - Analysis of Engineered Proteins by Fluorescence Quenching**

Acrylamide effectively quenches the fluorescence of accessible tryptophan residues in proteins. We examined fluorescence quenching of the single tryptophan residue of BHL1, BHL2, BHL3, BHL4, and wild-type CI-2 in the presence or absence of 6M guanidinium chloride. The quenching of intrinsic fluorescence of the proteins was followed by sequential addition of small aliquots of a 1 M acrylamide solution. The excitation wavelength was set at 295 nm to ensure optimal absorption by the tryptophan residue. In the absence of denaturant, an emission wavelength of 337 nm and a protein concentration of 20  $\mu$ M were used. In the presence of 6 M guanidinium chloride, the emission

wavelength was 356 nm and the protein concentration was lowered to 2  $\mu$ M because of the increase in the quantum yield of fluorescence after denaturation. The fluorescence intensities were corrected for the self-absorption of incident light [McClure and Edelman, Biochemistry 6: 567-572, (1967)) by using a molar extinction coefficient of 0.23 for acrylamide [Parker, "Photoluminescence of Solutions", Elsevier, New York, (1968)). The quenching data were plotted as a direct Stern-Volmer plot,  $F_0/F$  vs the molar concentration of acrylamide, where  $F_0$  is the fluorescence intensity in the absence of quencher and  $F$  is the fluorescence intensity in the presence of quencher. The Stern-Volmer quenching constant  $K_{sv}$  was determined from the slope of this plot, and is summarized in the following table.

*Stern-Volmer constants determined by acrylamide quenching of tryptophan fluorescence in the absence of denaturant (mean  $\pm$  standard deviation).*

Protein	$K_{sv}$ ( $M^{-1}$ )
WT CI-2	$1.7 \pm 0.1$
BHL1	$3.5 \pm 0.3$
BHL2, BHL3N	$5.5 \pm 0.4$
BHL3	$2.4 \pm 0.2$
BHL4	$0.65 \pm 0.02$

This experiment revealed that, in the absence of denaturant, there are differences in the accessibility of the tryptophan among these proteins. In contrast, the tryptophan was more completely accessible in all of the proteins upon unfolding in 6M guanidinium chloride (average  $K_{sv}$  of approximately 17  $M^{-1}$ ).

#### **Example 10 – Analysis of Engineered Proteins by Western Blots.**

Rabbit polyclonal antibodies (two rabbits for each) were prepared against truncated wild type CI-2, BHL1, or a 1:1 mixture of BHL6 and BHL8. Western blots of 100 ng of each protein were probed with a 1:1000 dilution of antisera against wild type CI-2 or against the BHL6/BHL8 mixture. Antisera to wild type CI-

2 reacted weakly with BHL5, BHL6, and BHL8, and reacted more strongly with all of the other BHL proteins and with wild type CI-2. Antibodies against the BHL6/BHL8 mixture reacted most strongly with BHL 5, BHL6, and BHL8, but reacted less strongly with the other BHL proteins and with wild-type CI-2. Other  
5 western blots revealed that antisera against BHL1 recognized wild type CI-2, BHL1, BHL2, BHL3, BHL3N, BHL4, BHL5, and BHL6.

### **Example 11 – Expression of Engineered Proteins in Plants**

Numerous constructs with various promoters and upstream and downstream regulatory elements have been prepared to express BHL8, BHL6, BHL4, and BHL3N in maize (corn), and plants have been transformed. BHL3N with a gamma zein promoter and with a heterologous signal peptide was expressed in corn endosperm, as demonstrated by positive western blots and ELISAs, using antibodies against BHL1. In contrast, the BHL3N protein expressed with the same promoter but with no signal peptide was not detected in transgenic corn, demonstrating that targeting this protein to the endoplasmic reticulum allowed higher expression than was possible with cytosolic (non-targeted) expression. In *Arabidopsis*, BHL5, BHL6, and BHL8 will be expressed with a constitutive promoter to further assess effects of protein stability on protein expression levels in plant leaves and seeds.

### **Example 12 – Fusion Proteins.**

A construct was prepared that encoded a BHL3N dimer, with one BHL3N molecule fused at the amino terminus to the carboxy terminus of the other BHL3N molecule. The BHL3N fusion protein was expressed in *E. coli* and purified. Fluorescence and circular dichroism analysis revealed conformational differences between the BHL3N fusion protein and the BHL3N monomer.

The BHL3N polypeptide could also be fused at its amino terminus through genetic engineering methods known in the art, to another protein enriched in essential amino acids, such as high lysine hordeothionin (Rao et al., Protein Engineering 7: 1485-1493, 1994). An amino terminal extension could also include a start signal, a transit sequence, a signal peptide, a fusion protein, a cleavable peptide, or an uncleaved peptide (we need examples and ref. here)

The amino terminus of the C1-2 derived protein may need to have the terminal methionine removed in order to ensure correct translation of the fusion polypeptide. It is known to one of skill in the art how to use restriction enzymes and oligonucleotides, to provide an intact nucleotide sequence that is in frame and able to be translated into the polypeptide of the invention.

### **Example 13 – Peptide insertions In Active Site Loop.**

It was previously shown that inserting peptides containing glutamine, alanine, or glycine in the active site loop region of wild type CI-2 had relatively minor effects on protein stability (Ladurner and Fersht, J. Mol. Biol. 273: 330-337, 1997). Peptides enriched in essential amino acids will be inserted into the active site loop region of the engineered proteins of the present study.

### **Example 14 – Substitutions**

The CI-2-like protein will be further modified by substituting one or more of the following: V32T; E45T, D64T, D74T, or A77T. Modifications will use materials and methods described supra utilizing any CI-2-like protein.

We claim:

1. An isolated polypeptide comprising a CI-2-like protein with a non-native essential amino acid residue in more than about 11% to less than about 80% of the amino acid residues.
2. The polypeptide of Claim 1 further comprising protein modification.
3. The polypeptide of Claim 1 further comprising from one to 5 disulfide-bonds.
4. The polypeptide of Claim 1 wherein the protein exhibits a free energy of unfolding of more than about 3.5 to less than about 15 Kilocalories per mole.
5. The polypeptide of Claim 1 wherein the polypeptide is proteolytically stable, as demonstrated by detection of the intact polypeptide based upon detection by SDS-PAGE analysis, following a 30 minute incubation at 37°C in 100mM Tris-HCl, 50mMNaCl, 1mM $\text{CaCl}_2$ , pH 8 , with a 10:1 (weight to weight ratio) of polypeptide:protease, with the protease being either chymotrypsin or trypsin.
6. The polypeptide of Claim 1 further comprising an amino-terminal extension.
7. The polypeptide of Claim 1 wherein the protein exhibits a modified protease activity.
8. The polypeptide of Claim 1 wherein the essential amino acid residues comprise lysine, threonine, tryptophan, methionine or combinations thereof.
9. An isolated polypeptide comprising a plant CI-2-like polypeptide altered to have the following composition: 15-35 mole % lysine, 5-15 mole % methionine, 6-25 mole % threonine, 4-9 mole % tryptophan or combinations thereof.
10. An isolated polypeptide comprising Seq. ID. No. 2, or truncated versions thereof, modified to contain 7 or more non-native essential amino acid residues at positions corresponding to the positions in Sequence ID. No. 2 selected from 1, 8, 11, 17, 18, 19, 20, 22, 23, 31, 34, 38, 40, 41, 47, 49, 56, 58, 59, 60, 61, 62, 63, 65, 67, 69, 73, 75, 76, 78, 79, 81, 82, or combinations thereof.
11. The polypeptide of Claim 10 wherein the essential amino acid residues comprise lysine, threonine, tryptophan, methionine, or combinations or conservative substitutions thereof.

12. The polypeptide of Claim 10 wherein the protein exhibits reduced inhibitory activity against chymotrypsin, subtilisin or elastase.

13. The protein of claim 10 wherein the polypeptide comprises one or more of the following modifications: V32T; E45T; D64T; D74T; or A77T.

5 14. The protein of claim 10 further comprising one of the following modifications: [T22C, V82C], [E23C, R81C] or [V53C, V70C].

15. The polypeptide of Claim 10 further comprising an amino-terminal extension.

16. The protein according to claim 15 wherein the amino terminal extension comprises a nutritionally-enhancing polypeptide.

10 17. The polypeptide of Claim 15 wherein the amino-terminal extension is a start signal, a transit sequence, a transit peptide, a signal peptide, a fusion protein, a cleavable peptide, a CI-2-like polypeptide or an uncleaved peptide.

18. The polypeptide of Claim 15 wherein the CI-2 derived polypeptide comprises at least 1 to about 18 additional residues corresponding to amino acid  
15 residues 1 to amino acid residue 17 of Seq. ID No. 2 or 12.

19. An isolated polypeptide comprising a CI-2 derived protein comprising two or more of the following modifications corresponding to positions in Seq. ID No. 2 selected from:

20 H18A, I, L, V or M; N19K or T; L20M I, or V; E23T or K; S31T or K; E34K or T; V38M I, or L; L40M I, or V; Q41K or T; Q47K or T; I49M I, L, or V; I56K or T; M59G; R62K or T; I63M, L, or V; R65K or T; R67K or T; F69W; L73K or T; A75K or T; Q78K or T; V79T or K; or R81K, or T.

20. The polypeptide according to claim 19 wherein the modifications comprise one or more of the following modifications: [E23C and R81C] or [T22C and  
25 V82C] or [V53C and V70C].

21. An isolated polypeptide comprising a CI-2 derived protein comprising two or more of the following modifications corresponding to positions in Seq. ID No. 2 selected from:

30 H18A or M; N19K; L20M; T22C; E23T or C; S31T; E34K; V38M; L40M; Q41K; Q47K; I49M; I56K; M59G; R62K; I63M; R65K; R67K; F69W; L73K; A75K; Q78K; V79T; R81K or C; or V82C.



22. The polypeptide of claim 21, further comprising substituting a tryptophan at position 61 and a glycine at position 59.

23. The polypeptide of claim 22, further comprising threonine at one or more of positions 32, 45, 53, 64 or 70.

5 24. The polypeptide according to claim 22 wherein the modifications comprise one or more of the following modifications: [E23C and R81C] or [T22C and V82C] or [V53C and V70C].

10 25. The polypeptide according to claim 22 further comprising an insert in the active site loop region that is enriched in essential amino acids for the purpose of nutritional enhancement.

26. An isolated polypeptide comprising a CI-2 derived protein comprising two or more of the following modifications corresponding to positions in Seq. ID No. 2 selected from:

15 [one or more of S1 or S2 or V3 or E4 or K5 or K6 or P7 or E8 or G9 or V10 or N11 or T12 or G13 or A14 or G15 or D16 deleted];  
S1K; E8K; N11K; [R17K or M]; [H18A or M]; N19K; L20M; T22 C; [E23T or C]; S31T; V32T; E34K; V38M; L40M; Q41K; E45T; Q47K; I49M; V53C;  
[[[I56K] and [T58A, or G] and [M59K or G] and [E60A or H] and [Y61W] and [R62K]] or [[I56K] and [M59K or G] and [Y61W] and [R62K]] or [[I56K] and [M59K or G] and [R62K]]];  
20 I63M; D64T; R65K; R67K; F69W; V70C; L73K; D74T; N75K; A77T; Q78K; V79T; [R81K or C]; or V82 C.

27. An isolated polypeptide comprising a CI-2 derived protein comprising modifications corresponding to positions in Seq. ID No. 2 selected from:

25 [[[I56K] and [T58A, or G] and [M59K or G] and [E60A or H] and [Y61W] and [R62K]] or  
[[[I56K] and [M59K or G] and [Y61W] and [R62K]] or [[I56K] and [M59K or G] and [R62K]]]; and two or more of the following modifications:  
[one or more of S1 or S2 or V3 or E4 or K5 or K6 or P7 or E8 or G9 or V10 or N11 or T12 or G13 or A14 or G15 or D16 deleted];  
30 S1K; E8K; N11K; R17K or M; [H18A or M]; N19K; L20M; T22 C; [E23T or C]; S31T; V32T; E34K; V38M; L40M; Q41K; E45T; Q47K; I49M; V53C;

I63M; D64T; R65K; R67K; F69W; V70C; L73K; D74T; N75K; A77T; Q78K; V79T; [R81K or C]; or V82 C.

28. An isolated polypeptide of Sequence ID. No. 2 comprising a protein with three or more non-native essential amino acids at positions selected from 1, 8, 11, 17, 18, 19, 20, 22, 23, 31, 32, 34, 38, 40, 41, 45, 47, 49, 56, 58, 59, 60, 61, 62, 63, 64, 65, 67, 69, 73, 74, 75, 76, 77, 78, 79, 81, 82, or combinations thereof and;

excluding V, P, W, S, E and R at position 56; S, K, R, P, E, V, Y, W, and A at position 58; R, Y, P, W, E, V, S, K, and A at position 59; Q, S, T, I, P, and K at position 60; V, E, R, P, and W at position 61 and E, Q, N, V, F, and Y position 62 and

conservatively modified and conservatively substituted variants thereof.

29. An isolated polypeptide comprising Seq. ID No. 6, 8, 10, 12, 14, 16, 18, 20 or conservatively modified or conservatively substituted variants thereof.

30. An isolated polypeptide comprising at least 23 contiguous amino acids of SEQ. ID Nos. 6, 8, 10, 12, 14, 16, 18, 20.

31. An isolated polypeptide comprising at least 23 contiguous amino acids with more than 79% sequence identity, to the polypeptide of Seq. ID No. 20, wherein the % sequence identity is based on the 23 contiguous amino acids sequence and is determined by GAP analysis using Gap Weight of 12 and Length Weight of 4.

32. An isolated polypeptide that is immunologically reactive with antibodies against the protein of Seq. ID No. 20 and not SEQ ID No. 2.

33. An isolated nucleic acid comprising:

(a) a polynucleotide encoding the protein of claim 1;

(b) a polynucleotide that encodes a polypeptide of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, or 20;

(c) a polynucleotide amplified from a plant nucleic acid library using the primers of SEQ ID NOS: 21 and 22;

(d) a polynucleotide comprising at least 20 contiguous bases of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17 or 19;

- (e) a polynucleotide encoding a plant CI-2-derived polypeptide having 15% more essential amino acids than SEQ ID NO 2;
- (f) a polynucleotide having at least 73% sequence identity to SEQ ID NO: 19, wherein the % sequence identity is based on the entire sequence and is determined by BLAST 2.0;
- (g) a polynucleotide comprising at least 25 nucleotides in length which hybridizes under low stringency conditions to a polynucleotide having the sequence set forth in SEQ ID NOs: 19, wherein the conditions include hybridization with a buffer solution of 30% formamide, 1 M NaCl, 1% SDS at 37°C for 24 hours and a wash in 2X SSC at 50°C, 3x for 15 minutes;
- (h) a polynucleotide comprising the sequence set forth in SEQ ID NOs: 5, 7, 9, 11, 13, 15, 17 or 19;
- (i) conservatively modified variants of SEQ ID NO : 5, 7, 9, 11, 13, 15, 17 or 19; or
- (j) a polynucleotide complementary to a polynucleotide of (a) through (i).

34. The isolated nucleic acid of claim 33 wherein the polynucleotide is a plant polynucleotide.

35. A vector comprising at least one nucleic acid of claim 33.

36. An expression cassette comprising at least one nucleic acid of claim 33 operably linked to a promoter, wherein the nucleic acid is in sense or antisense orientation.

37. A host cell into which is introduced at least one expression cassette of claim 36.

38. The host cell of claim 37 that is a plant cell.

39. A transgenic plant comprising at least one expression cassette of claim 36.

40. The transgenic plant of claim 39, wherein the plant is corn, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, lupin or millet.

41. A seed from the transgenic plant of claim 40.

42. The seed of claim 41, wherein the seed is from corn, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, lupin or millet.

43. A ribonucleic acid sequence encoding a polypeptide of claim 10.

44. A method for increasing the essential amino acid content in a polypeptide in a plant, comprising:

- (a) stably transforming a plant cell with the polynucleotide encoding the polypeptide of claim 10, operably linked to a promoter, wherein the polynucleotide is in sense orientation;
- (b) growing the plant cell under plant growing conditions to produce a regenerated plant; and
- (c) expressing the polypeptide for a time sufficient to produce the polypeptide encoded by the polynucleotide of (a) in the plant.

45. The method of claim 44, wherein the plant cell is corn, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, lupin or millet.

46. A method of increasing expression levels of a protein in a transgenic plant comprising:

- engineering a nucleotide sequence encoding the protein of interest to increase the in-vitro proteolytic or thermodynamic stability of the protein;
- introducing at least 1 copy into a plant cell; and
- expressing the protein.

47. A method of increasing nutritional value of feed comprising substituting more than 11% to less than 75% of the amino acids residues of a protein with essential amino acids and modifying the protein to increase the stability of the in-vivo expressed polypeptide.

48. The method of Claim 47 wherein the protein is a CI-2-like polypeptide.

49. The method of Claim 47 wherein the modifying stability is from one or more disulfide bonds.

50. A method of increasing nutritional value of a protein by altering a CI-2 homologue to enhance its nutritional value by altering amino acid residues to the positions in Sequence ID. No. 2 selected from 1, 8, 11, 17, 18, 19, 20, 22, 23, 31, 34, 38, 40, 41, 47, 49, 56, 58, 59, 60, 61, 62, 63, 65, 67, 69, 73, 75, 76, 78, 79, 81, 82, or combinations thereof.

51. An isolated nucleic acid comprising:

- (a) a polynucleotide of Seq ID Nos 23, 25, 27, 29 and 31;

- (b) a polynucleotide that encodes a polypeptide of SEQ ID NOS: 24, 26, 28, 30, 32;
- (c) a polynucleotide amplified from a Zea mays nucleic acid library using the primers of SEQ ID NOS: 21 and 22;
- 5 (d) a polynucleotide comprising at least 20 contiguous bases of SEQ ID NOS: 23, 25, 27, 29 and 31;
- (e) a polynucleotide having at least 50% sequence identity to SEQ ID NOS: 23, 25, 27, 29 and 31, wherein the % sequence identity is based on the entire sequence and is determined by BLAST 2.0;
- 10 (f) a polynucleotide comprising at least 25 nucleotides in length which hybridizes under low stringency conditions to a polynucleotide having the sequence set forth in SEQ ID NOS: 23, 25, 27, 29 and 31, wherein the conditions include hybridization with a buffer solution of 30 % formamide, 1 M NaCl, 1% SDS at 37°C for 4-12 hours and a wash in 2X SSC at 50°C;
- 15 (g) a polynucleotide comprising the sequence set forth in SEQ ID NOS: 23, 25, 27, 29 and 31;
- (h) conservatively modified variants of SEQ ID NO 23, 25, 27, 29 and 31; or
- 20 (i) a polynucleotide complementary to a polynucleotide of (a) through (h).

25

\_\_\_\_\_

## ABSTRACT OF THE DISCLOSURE

The invention provides isolated nucleic acids and their encoded polypeptides that are involved in enhancing the essential amino acid content of a plant. Optionally there is also a decrease in protease inhibitory activity of the polypeptide. The invention further provides recombinant expression cassettes, host cells, transgenic plants, and antibody compositions. The present invention provides methods and compositions relating to increasing essential amino acid content of plants for feed.

Figure 1

SEQUENCES OF BHL PROTEINS

	10	20	30	40	50	60	70	80
	*	*	*	*	*	*	*	*
WT CI - 2 (native)	SSVEKKPEGVNTGAGDRHNLKTEWPELVGKSVEAKKVIILQDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
WT CI-2	NMLKTEWPELVGKSVEAKKVIILQDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL1	MKLKTEWPELVGKSVEAKKVIILKDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL2	MKLKTEWPELVGKSVEAKKVIILKDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL3	MKLKTEWPELVGKSVEAKKVIILKDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL3N	MKSVEKKPKGVKTGAGDKHKLKTEWPELVGKSVEAKKVIILKDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL4	MKLKTEWPELVGKSVEAKKVIILKDKPEAQIIVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL5	MAKMKTTWPELVGKIVEAKKIMKDKPEAKIMVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL6	MAKMKTTWPELVGKIVEAKKIMKDKPEAKIMVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHL8	MAKMKCTWPELVGKIVEAKKIMKDKPEAKIMVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHLX	XXXXXXXXXXWPELVGKIVEAKKIXXDKPEAXIXVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							
BHLXP	XXXXXXXXXXWPELVGKIVEAKKIXXDKPEAXIXVLPVGTIVTMEYRIDRVRLFVDKLDNIAQVPRVG							

	20	30	40	50	60	70	80
	*	*	*	*	*	*	*
WT CI-2	MNLKTEWPELVGKSVEEAKKVLQDKPEAQIIVLPV-GTIV-TMEYRIDRVRLFVDKLDNIAQVPRVG						
Homologs							
1	GAKTSWPEVVGMSAEKAKEIILRDKPNAQIEVIPV-DAMV-PLNFNPNRVFLVHKATTVAZVSRVG						
2	GAKTEWPELVGCTIKEAKEKIKADRPDLKVIVP-GSIV-TQEIIDLNRRVRVWDKV--AKVPKIG						
3	GAKTSWPEVVGSLVEDAKKVLKDKPDADIVLPV-GSVV-TADYRPNRVRIEVDIV--AQTPHIG						
4	--RTSWPELVGVSAAEEARK-IKEEMPEAEIQVVPQ-DSFV-TADYKFQRVRLYVDESNKVVRAAPIG						
5	AEKSSWPELVGEDGEEAVKIIQENPSLDVILMPR-GQNWATLDCRPNRVRVFNDESGKVN SIPRIG						
6	TRKTSWPELVGTAAEEAE-KIKEEMSGVEIQVVP-GSFV-TADYKPKQRVRLYVDESNKVT RTPGIG						
7	PTKTSWPELVGVTAEQAETKIKEEMVDVQIQVSPH-DSFV-TADYNPKRVRKYVDESNKVT RTPSIG						
8	VTKERWPELLGTPAKFAMQIIQENPKLTNVQTVLNGTPV-TEDLRCNRVRLFVNVLDFVVQTPQVG						
9	NKKETWPELIGVPAKFAREIIQENSKLTNP SVLNGSPV-TKDFRCERVRRLFVNVLDFVVQIPRVG						
10	VGKNTWPELTCGARGEEAATVETENP SVTAVIPE-GSIV-TTDERCDRVRVWVDENGIVTRVPVIG						
11	QGKSSWPQLVGSTGAATAKAVIERENPVRRAVILKV-GSGA-TKDFRCDRVRVWVTERGIVARPTIG						
12	PGKSSWPHLVGGGSVAKAIIERQNPVKAVIL EE-GTPV-TKDFRCNRVRLIWNKRGLVVSPPRIG						
13	DGKLQWPELIGVPTKLAKELIEKQNSLISNVHILLNGSPV-TMDFRCNRVRLFDDILG SVVQIPRVA						
14	NGKLSWPELIGVPAHYAKGII EKENS LITNVQIILLNGSPV-TMDYRCNRVRLF DNILG DVVQIPRVA						
15	KGKQFWPELIGVPALYAKGII EKENS ITNIPILLNGSPV-TKDFRCDRVRRLFVNILG DVVQIPRV						
16	VTKESWPELLGTPAKFAKQIIQENPKLTNVEELLNGSAF-TEDLRCNRVRRLFVNLLDIVVQTPKV						
17	EGKQMPPELIGVPTKLAKELIEKENPSITNIPILLSGSPI-TLDYLCDRVRRLF DNILG FVVQMPVVT						
18	PGKQEWPELVGEYGYKAAAI IERENPNVRSIVKHE-RSGF-TKDFRCDRVWVVDSTGVVVRTPRVT						
19	VIFNSWSVLGTNGDYAAVVERENPTVNAAVILLD-GSPV-TADFCRCDRVRVFVDGNRI VVKTPKSG						



## SEQUENCE LISTING

<110> Pioneer Hi-Bred International, Inc.

<120> Proteins With Enhanced Levels of  
Essential Amino Acids

<130> 0571R2

<150> 08/740,682

<151> 1996-11-01

<150> PCT/US97/20441

<151> 1997-10-31

<160> 32

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 249

<212> DNA

<213> Hordeum vulgare

<220>

<221> CDS

<222> (1) ... (249)

<400> 1

agt tca gtg gag aag aag ccg gag gga gtg aac acc ggt gct ggt gac	48
Ser Ser Val Glu Lys Lys Pro Glu Gly Val Asn Thr Gly Ala Gly Asp	
1 5 10 15	

cgt cac aac ctg aag aca gag tgg cca gag ttg gtg ggg aaa tcg gtg	96
Arg His Asn Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val	
20 25 30	

gag gag gcc aag aag gtg att ctg cag gac aag cca gag gcg caa atc	144
Glu Glu Ala Lys Lys Val Ile Leu Gln Asp Lys Pro Glu Ala Gln Ile	
35 40 45	

ata gtt cta ccg gtg ggg aca att gtg acc atg gaa tat cgg atc gac	192
Ile Val Leu Pro Val Gly Thr Ile Val Thr Met Glu Tyr Arg Ile Asp	
50 55 60	

cgc gtc cgc ctc ttt gtc gat aaa ctc gac aac att gcc cag gtc ccc	240
Arg Val Arg Leu Phe Val Asp Lys Leu Asp Asn Ile Ala Gln Val Pro	
65 70 75 80	

agg gtc ggc	249
Arg Val Gly	

<210> 2

<211> 83

<212> PRT

<213> Hordeum vulgare

<400> 2

662730-663460

<210> 5

<211> 198  
 <212> DNA  
 <213> Hordeum vulgare

<220>  
 <221> CDS  
 <222> (1)...(198)

<400> 5

atg aag ctg aag aca gag tgg ccg gag ttg gtg ggg aaa tcg gtg gag 48  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15

aaa gcc aag aag gtg atc ctg aag gac aag cca gag gcg caa atc ata 96  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30

gtt ctg ccg gtt ggt aca aag gtg acg aag gaa tat aag atc gac cgc 144  
 Val Leu Pro Val Gly Thr Lys Val Thr Lys Glu Tyr Lys Ile Asp Arg  
 35 40 45

gtc aag ctc ttt gtg gat aaa aag gac aac atc gcg cag gtc ccc agg 192  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60

gtc ggc 198  
 Val Gly  
 65

<210> 6  
 <211> 66  
 <212> PRT  
 <213> Hordeum vulgare

<400> 6

Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30  
 Val Leu Pro Val Gly Thr Lys Val Thr Lys Glu Tyr Lys Ile Asp Arg  
 35 40 45  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60  
 Val Gly  
 65

<210> 7  
 <211> 198  
 <212> DNA  
 <213> Hordeum vulgare

<220>  
 <221> CDS  
 <222> (1)...(198)

<400> 7

atg aag ctg aag aca gag tgg ccg gag ttg gtg ggg aaa tcg gtg gag 48  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15

aaa gcc aag aag gtg atc ctg aag gac aag cca gag gcg caa atc ata 96

66250-66250

Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30

gtt cta ccg gtt ggt aca aag gtg gcg aag gcc tat aag atc gac aag 144  
 Val Leu Pro Val Gly Thr Lys Val Ala Lys Ala Tyr Lys Ile Asp Lys  
 35 40 45

gtc aag ctt ttt gtg gat aaa aag gac aac atc gcg cag gtc ccc agg 192  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60

gtc ggc 198  
 Val Gly  
 65

<210> 8

<211> 66

<212> PRT

<213> Hordeum vulgare

<400> 8

Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30  
 Val Leu Pro Val Gly Thr Lys Val Ala Lys Ala Tyr Lys Ile Asp Lys  
 35 40 45  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60  
 Val Gly  
 65

<210> 9

<211> 198

<212> DNA

<213> Hordeum vulgare

<220>

<221> CDS

<222> (1)...(198)

<400> 9

atg aag ctg aag aca gag tgg ccg gag ttg gtg ggg aaa tcg gtg gag 48  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15

aaa gcc aag aag gtg atc ctg aag gac aag cca gag gcg caa atc ata 96  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30

gtt cta ccg gtt ggt aca aag gtg ggt aag cat tat aag atc gac aag 144  
 Val Leu Pro Val Gly Thr Lys Val Gly Lys His Tyr Lys Ile Asp Lys  
 35 40 45

gtc aag ctt ttt gtg gat aaa aag gac aac atc gcg cag gtc ccc agg 192  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60

gtc ggc 198  
 Val Gly  
 65

66T50"669T50

<210> 10  
 <211> 66  
 <212> PRT  
 <213> Hordeum vulgare

<400> 10  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
 1 5 10 15  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
 20 25 30  
 Val Leu Pro Val Gly Thr Lys Val Gly Lys His Tyr Lys Ile Asp Lys  
 35 40 45  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
 50 55 60  
 Val Gly  
 65

<210> 11  
 <211> 252  
 <212> DNA  
 <213> Hordeum vulgare

<220>  
 <221> CDS  
 <222> (1)...(252)

<400> 11  
 atg aag tcg gtg gag aag aaa ccg aag ggt gtg aag aca ggt gcg ggt 48  
 Met Lys Ser Val Glu Lys Lys Pro Lys Gly Val Lys Thr Gly Ala Gly  
 1 5 10 15  
 gac aag cat aag ctg aag aca gag tgg ccg gag ttg gtg ggg aaa tcg 96  
 Asp Lys His Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser  
 20 25 30  
 gtg gag aaa gcc aag aag gtg atc ctg aag gac aag cca gag gcg caa 144  
 Val Glu Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln  
 35 40 45  
 atc ata gtt cta ccg gtt ggt aca aag gtg ggt aag cat tat aag atc 192  
 Ile Ile Val Leu Pro Val Gly Thr Lys Val Gly Lys His Tyr Lys Ile  
 50 55 60  
 gac aag gtc aag ctt ttt gtg gat aaa aag gac aac atc gcg cag gtc 240  
 Asp Lys Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val  
 65 70 75 80  
 ccc agg gtc ggc 252  
 Pro Arg Val Gly

<210> 12  
 <211> 84  
 <212> PRT  
 <213> Hordeum vulgare

<400> 12  
 Met Lys Ser Val Glu Lys Lys Pro Lys Gly Val Lys Thr Gly Ala Gly  
 1 5 10 15

66250-531-666

Asp Lys His Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser  
                   20                  25                  30  
 Val Glu Lys Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln  
                   35                  40                  45  
 Ile Ile Val Leu Pro Val Gly Thr Lys Val Gly Lys His Tyr Lys Ile  
                   50                  55                  60  
 Asp Lys Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val  
 65                  70                  75                  80  
 Pro Arg Val Gly

<210> 13  
 <211> 198  
 <212> DNA  
 <213> Hordeum vulgare

<220>  
 <221> CDS  
 <222> (1)...(198)

<400> 13  
 atg aag ctg aag aca gag tgg ccg gag ttg gtg ggg aaa tcg gtg gag 48  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
   1                  5                  10                  15  
 aaa gcc aag aag gtg atc ctg aag gac aag cca gag gcg caa atc ata 96  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
                   20                  25                  30  
 gtt cta ccg gtt ggt aca aag gtg acg ggc gaa tac aag atc gac cgc 144  
 Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Tyr Lys Ile Asp Arg  
                   35                  40                  45  
 gtc aag ctt ttt gtg gat aaa aag gac aac atc gcg cag gtc ccc agg 192  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
                   50                  55                  60  
 gtc ggc 198  
 Val Gly  
 65

<210> 14  
 <211> 66  
 <212> PRT  
 <213> Hordeum vulgare

<400> 14  
 Met Lys Leu Lys Thr Glu Trp Pro Glu Leu Val Gly Lys Ser Val Glu  
   1                  5                  10                  15  
 Lys Ala Lys Lys Val Ile Leu Lys Asp Lys Pro Glu Ala Gln Ile Ile  
                   20                  25                  30  
 Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Tyr Lys Ile Asp Arg  
                   35                  40                  45  
 Val Lys Leu Phe Val Asp Lys Lys Asp Asn Ile Ala Gln Val Pro Arg  
                   50                  55                  60  
 Val Gly  
 65

<210> 15  
 <211> 201  
 <212> DNA

662F50"669F660

<213> Hordeum vulgare

<220>

<221> CDS

<222> (1) ... (201)

<400> 15

atg gct aag atg aag aca acg tgg cct gag ctg gtg ggc aag acc gtg	48
Met Ala Lys Met Lys Thr Thr Trp Pro Glu Leu Val Gly Lys Thr Val	
1 5 10 15	

gag aaa gcc aag aag atg atc atg aag gac aag cca gag gcg aag atc	96
Glu Lys Ala Lys Lys Met Ile Met Lys Asp Lys Pro Glu Ala Lys Ile	
20 25 30	

atg gtt ctg cca gtt ggg acc aaa gtg acc ggt gaa tgg aag atg gat	144
Met Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Trp Lys Met Asp	
35 40 45	

cgc gtc aaa ctc tgg gtc gac aag aag gac aag atc gcc aag act ccg	192
Arg Val Lys Leu Trp Val Asp Lys Lys Asp Lys Ile Ala Lys Thr Pro	
50 55 60	

aag gtc ggc	201
Lys Val Gly	
65	

<210> 16

<211> 67

<212> PRT

<213> Hordeum vulgare

<400> 16

Met Ala Lys Met Lys Thr Thr Trp Pro Glu Leu Val Gly Lys Thr Val	
1 5 10 15	
Glu Lys Ala Lys Lys Met Ile Met Lys Asp Lys Pro Glu Ala Lys Ile	
20 25 30	
Met Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Trp Lys Met Asp	
35 40 45	
Arg Val Lys Leu Trp Val Asp Lys Lys Asp Lys Ile Ala Lys Thr Pro	
50 55 60	
Lys Val Gly	
65	

<210> 17

<211> 201

<212> DNA

<213> Hordeum vulgare

<220>

<221> CDS

<222> (1) ... (201)

<400> 17

atg gct aag atg aag aca acg tgg cct gag ctg gtg ggc aag acc gtg	48
Met Ala Lys Met Lys Thr Thr Trp Pro Glu Leu Val Gly Lys Thr Val	
1 5 10 15	

gag aaa gcc aag aag atg atc atg aag gac aag cca gag gcg aag atc	96
Glu Lys Ala Lys Lys Met Ile Met Lys Asp Lys Pro Glu Ala Lys Ile	
20 25 30	

66250"6694450

atg gtt ctg cca gtt ggg acc aaa gtg acc ggt gaa tgg aag atg gat 144  
Met Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Trp Lys Met Asp  
35 40 45

cgc gtc cgc ctc tgg gtc gac aag aag gac aag atc gcc aag act ccg 192  
Arg Val Arg Leu Trp Val Asp Lys Lys Asp Lys Ile Ala Lys Thr Pro  
50 55 60

aag gtc ggc 201  
Lys Val Gly  
65

<210> 18  
<211> 67  
<212> PRT  
<213> Hordeum vulgare

<400> 18  
Met Ala Lys Met Lys Thr Thr Trp Pro Glu Leu Val Gly Lys Thr Val  
1 5 10 15  
Glu Lys Ala Lys Lys Met Ile Met Lys Asp Lys Pro Glu Ala Lys Ile  
20 25 30  
Met Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Trp Lys Met Asp  
35 40 45  
Arg Val Arg Leu Trp Val Asp Lys Lys Asp Lys Ile Ala Lys Thr Pro  
50 55 60  
Lys Val Gly  
65

<210> 19  
<211> 201  
<212> DNA  
<213> Hordeum vulgare

<220>  
<221> CDS  
<222> (1)...(201)

<400> 19  
atg gct aag atg aag tgc acg tgg cct gag ctg gtg ggc aag acc gtg 48  
Met Ala Lys Met Lys Cys Thr Trp Pro Glu Leu Val Gly Lys Thr Val  
1 5 10 15

gag aaa gcc aag aag atg atc atg aag gac aag cca gag gcg aag atc 96  
Glu Lys Ala Lys Lys Met Ile Met Lys Asp Lys Pro Glu Ala Lys Ile  
20 25 30

atg gtt ctg cca gtt ggg acc aaa gtg acc ggt gaa tgg aag atg gat 144  
Met Val Leu Pro Val Gly Thr Lys Val Thr Gly Glu Trp Lys Met Asp  
35 40 45

cgc gtc cgc ctc tgg gtc gac aag aag gac aag atc gcc aag act ccg 192  
Arg Val Arg Leu Trp Val Asp Lys Lys Asp Lys Ile Ala Lys Thr Pro  
50 55 60

aag tgc ggc 201  
Lys Cys Gly  
65

55CT50" 689T660





ggc ggc ggc gcc aag acg tcg tgg cct gag gtg gtc ggg ctg agc gtg 144  
 Gly Gly Gly Ala Lys Thr Ser Trp Pro Glu Val Val Gly Leu Ser Val  
 35 40 45  
 gag gac gcc aag aag gtg atg gtc aag gac aag ccg gac gcc gac atc 192  
 Glu Asp Ala Lys Lys Val Met Val Lys Asp Lys Pro Asp Ala Asp Ile  
 50 55 60  
 gtg gtg ctg ccc gtc ggc tcc gtg gtg acc gcg gat tat cgc cct aac 240  
 Val Val Leu Pro Val Gly Ser Val Val Thr Ala Asp Tyr Arg Pro Asn  
 65 70 75 80  
 cgt gtc cgc atc ttc gtc gac atc gtc gcc cag acg ccc cac atc ggc 288  
 Arg Val Arg Ile Phe Val Asp Ile Val Ala Gln Thr Pro His Ile Gly  
 85 90 95  
 tgataatata taagctagcc gctatttccct ttccttgccc cagaacttga aataaatata 348  
 tatacgatga aataacgcgg gcatgccgaa tanatggant gtgnntgaat tctcactaat 408  
 taagtaatgn cataaataaa cgtattcaaa aaaaaaaaaa aaaaaaaaaa a 459

<210> 24  
 <211> 96  
 <212> PRT  
 <213> Zea mays  
 <220>  
 <221> VARIANT  
 <222> (1)...(146)  
 <223> Xaa = Any Amino Acid

<400> 24  
 Ala Val Gln Gln Ala Arg Phe Thr Cys Pro Ser Ile Ile Ser Ser Thr  
 1 5 10 15  
 Gly Pro Ala Val Arg Asp Thr Met Ser Ser Thr Glu Cys Gly Gly Gly  
 20 25 30  
 Gly Gly Gly Ala Lys Thr Ser Trp Pro Glu Val Val Gly Leu Ser Val  
 35 40 45  
 Glu Asp Ala Lys Lys Val Met Val Lys Asp Lys Pro Asp Ala Asp Ile  
 50 55 60  
 Val Val Leu Pro Val Gly Ser Val Val Thr Ala Asp Tyr Arg Pro Asn  
 65 70 75 80  
 Arg Val Arg Ile Phe Val Asp Ile Val Ala Gln Thr Pro His Ile Gly  
 85 90 95

<210> 25  
 <211> 428  
 <212> DNA  
 <213> Zea mays

<220>  
 <221> CDS  
 <222> (1)...(303)

<400> 25  
 cga ccc acg cgt ccg ccc acg cgt ccg gca aga ttt acc tgc cca tcg 48  
 Arg Pro Thr Arg Pro Pro Thr Arg Pro Ala Arg Phe Thr Cys Pro Ser  
 1 5 10 15  
 atc ata tcg tca act ggt ccg gca gtt cgc gac acc atg agc tcc acg 96  
 Ile Ile Ser Ser Thr Gly Pro Ala Val Arg Asp Thr Met Ser Ser Thr  
 20 25 30

0311699.051399

```

<210> 26
<211> 101
<212> PRT
<213> Zea mays

<400> 26
Arg Pro Thr Arg Pro Thr Arg Pro Ala Arg Phe Thr Cys Pro Ser
 1          5          10          15
Ile Ile Ser Ser Thr Gly Pro Ala Val Arg Asp Thr Met Ser Thr
          20          25          30
Glu Cys Gly Gly Gly Gly Gly Gly Ala Lys Thr Ser Trp Pro Glu Val
          35          40          45
Val Gly Leu Ser Val Glu Asp Ala Lys Lys Val Ile Leu Lys Asp Lys
          50          55          60
Pro Asp Ala Asp Ile Val Val Leu Pro Val Gly Ser Val Val Thr Ala
65          70          75          80
Asp Tyr Arg Pro Asn Arg Val Arg Ile Phe Val Asp Ile Val Ala Gln
          85          90          95
Thr Pro His Ile Gly
          100

```

```
<220>
<221> CDS
<222> (1)...(255)

<221> misc_feature
<222> (1)...(441)
<223> n = A,T,C or G
```

<400> 27  
tta att att gcc ctt tca gtt ngc cat cgg cag ccg agc acc atg agc 48  
Leu Ile Ile Ala Leu Ser Val Xaa His Arg Gln Pro Ser Thr Met Ser  
1 5 10 15

<400> 29

gtg cgt cgt cgg cga aca gcc acc ggc ggc aag acg tcg tgg ccg gag 48  
 Val Arg Arg Arg Arg Thr Ala Thr Gly Gly Lys Thr Ser Trp Pro Glu  
 1 5 10 15

gtg gtc ggg ctg agc gtc gag gaa gcc aag aag gtg att ctg gcg gac 96  
 Val Val Gly Leu Ser Val Glu Glu Ala Lys Lys Val Ile Leu Ala Asp  
 20 25 30

aag ccg aac gcc gac atc gtg gtg ctg ccc acc acc acg cag gcg gtg 144  
 Lys Pro Asn Ala Asp Ile Val Val Leu Pro Thr Thr Thr Gln Ala Val  
 35 40 45

acc tcc gac ttt ggg ttc gac cgt gtc cgc gtc ttc gtc ggg acc gtc 192  
 Thr Ser Asp Phe Gly Phe Asp Arg Val Arg Val Phe Val Gly Thr Val  
 50 55 60

gcc cag acg ccc cat gtt ggc taggctagag cctcagccta gaggtcgctcg 243  
 Ala Gln Thr Pro His Val Gly  
 65 70

gcaccgccgg ccatgaccac ctgctantat gtcactnact agtaataaag tatwaataac 303  
 agggaggatg catgctcatc nttggaatct gtacgcttgt tggactacta cttggctact 363  
 tgaaaaaaaa aaaaaaaaaa 382

<210> 30  
 <211> 71  
 <212> PRT  
 <213> Zea mays

<400> 30  
 Val Arg Arg Arg Arg Thr Ala Thr Gly Gly Lys Thr Ser Trp Pro Glu  
 1 5 10 15  
 Val Val Gly Leu Ser Val Glu Glu Ala Lys Lys Val Ile Leu Ala Asp  
 20 25 30  
 Lys Pro Asn Ala Asp Ile Val Val Leu Pro Thr Thr Thr Gln Ala Val  
 35 40 45  
 Thr Ser Asp Phe Gly Phe Asp Arg Val Arg Val Phe Val Gly Thr Val  
 50 55 60  
 Ala Gln Thr Pro His Val Gly  
 65 70

<210> 31  
 <211> 448  
 <212> DNA  
 <213> Zea mays

<220>  
 <221> CDS  
 <222> (1)...(240)  
 <221> misc\_feature  
 <222> (1)...(448)  
 <223> n = A,T,C or G

<400> 31  
 cga ttt agc tat agc agg tct cga tcg gcg gcc atg agc ggt agc cgc 48  
 Arg Phe Ser Tyr Ser Arg Ser Arg Ser Ala Ala Met Ser Gly Ser Arg  
 1 5 10 15

agc aag aag tcg tgg ccg gag gtg gag ggg ctg ccg tcc gag gtg gcc 96  
 Ser Lys Lys Ser Trp Pro Glu Val Glu Gly Leu Pro Ser Glu Val Ala  
 20 25 30

66250-66345

```

aag cag aaa att ctg gcc gac cgc ccg gac gtc cag gtg gtc gtt ctg      144
Lys Gln Lys Ile Leu Ala Asp Arg Pro Asp Val Gln Val Val Val Leu
      35                      40                      45

ccc gac ggc tcc ttc gtc acc act gat ttc aac gac aag cgc gtc cgg      192
Pro Asp Gly Ser Phe Val Thr Thr Asp Phe Asn Asp Lys Arg Val Arg
      50                      55                      60

gtc ttc gtc gac aac gcc gac aac gtc gcc aaa gtc ccc aag atc ggc      240
Val Phe Val Asp Asn Ala Asp Asn Val Ala Lys Val Pro Lys Ile Gly
      65                      70                      75                      80

tagctagcta gctaggccca atcggttctaa tcagctagtt tctttctttc ataaataaaaa      300
gtcctctctc gtaccccgac tgtgatgttt ccctagttgt ctggtacgtg ttgttttctg      360
tcttaatgga tgccatggcg ccgcgcgcgc cctycatcat gaaaagctac atttgaaacg      420
attnnagta ttctttgctg ttaaaaaa      448

```

```

<210> 32
<211> 80
<212> PRT
<213> Zea mays

```

```

<400> 32
Arg Phe Ser Tyr Ser Arg Ser Arg Ser Ala Ala Met Ser Gly Ser Arg
 1          5          10          15
Ser Lys Lys Ser Trp Pro Glu Val Glu Gly Leu Pro Ser Glu Val Ala
      20          25          30
Lys Gln Lys Ile Leu Ala Asp Arg Pro Asp Val Gln Val Val Val Leu
      35          40          45
Pro Asp Gly Ser Phe Val Thr Thr Asp Phe Asn Asp Lys Arg Val Arg
      50          55          60
Val Phe Val Asp Asn Ala Asp Asn Val Ala Lys Val Pro Lys Ile Gly
      65          70          75          80

```

CCCTG"689F650